

Equilibrium sequences of irrotational binary polytropic stars : The case of double polytropic stars

Keisuke Taniguchi* and Takashi Nakamura**

* *Département d'Astrophysique Relativiste et de Cosmologie, UPR 8629 du C.N.R.S., Observatoire de Paris, F-92195 Meudon Cedex, France*

** *Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-8502, Japan*
(February 1, 2008)

Solutions to equilibrium sequences of irrotational binary polytropic stars in Newtonian gravity are expanded in a power of $\epsilon = a_0/R$, where R and a_0 are the orbital separation of the binary system and the radius of each star for $R = \infty$. For each order of ϵ , we should solve ordinary differential equations for arbitrary polytropic indices n . We show solutions for polytropic indices $n = 0.5, 1, 1.5$ and 2 up to ϵ^6 orders. Our semi-analytic solutions can be used to check the validity of numerical solutions.

PACS number(s): 04.30.Db, 04.25.Dm, 97.80.Fk

I. INTRODUCTION

Coalescing binary neutron stars (BNSs) are considered to be one of the most promising sources of gravitational waves for laser interferometers such as TAMA300 [1], GEO600 [2], VIRGO [3] and LIGO [4,5]. If we have accurate theoretical templates of inspiraling phase of BNSs, we can determine the mass and the spin of neutron stars from the gravitational wave signals in the inspiraling phase [6]. We may also extract the informations on the equation of state of a neutron star from the signals in the premerging phase [7]. For this purpose it is important to complete theoretical templates of gravitational waves in the premerging phase as well as in the inspiraling phase. Moreover, the study of BNSs in the premerging phase is motivated by the need to provide the realistic initial condition for the merging phase simulations [8–10].

In order to obtain accurate theoretical templates of gravitational waves from BNSs around the premerging phase [11], a general relativistic hydrostatic problem with compressible equation of state must be numerically solved. In solving this problem, we suppose that BNSs reach quasi-equilibrium states because the timescale of the orbital decay driven by the radiation reaction is much longer than the orbital period of BNSs until their innermost stable circular orbit (ISCO). (The formalism for solving quasi-equilibrium figures of irrotational binary neutron stars in general relativity is given in the references [12–15].) Furthermore, we regard the internal state of a neutron star as an irrotational or nearly irrotational one because the viscosity of a neutron star is not large enough for synchronization even near the ISCO [16,17].

Recently Bonazzola, Gourgoulhon and Marck have numerically studied the irrotational binary neutron stars in general relativity with the conformally flat condition [18–20]. More recently, Uryū and Eriguchi have also numerically solved the same problem as that of Bonazzola, Gourgoulhon and Marck [21]. In order to check the validity of these numerical calculations, it is necessary to compare them with analytic or semi-analytic ones. However, we have not obtained such analytic solutions in general relativity yet. There are solutions of irrotational BNSs only in the first post-Newtonian (1PN) approximation of general relativity. For example, Lombardi, Rasio and Shapiro have semi-analytically studied irrotational BNSs by using the energy variational method [22], and one of the authors of the present paper (KT) has analytically calculated equilibrium sequences of irrotational BNSs by using the tensor virial method [23]. The former one restricts the internal motion within the plane orthogonal to the rotational axis assuming the shape of the star to an ellipsoidal one in order to treat compressible equation of state. While in the latter one the velocity component along the orbital axis is included at 1PN order and the shape of the star at 1PN order is not restricted to an ellipsoidal one although the fluid is incompressible.

The situation is also the same in Newtonian gravity, that is, even in Newtonian gravity useful analytic or semi-analytic solutions for equilibrium sequences of irrotational binary systems are not obtained. Let us consider numerically constructed stationary structures of irrotational binary stars computed by Uryū and Eriguchi in Newtonian gravity [24,25]. One may think that semi-analytic solutions by Lai, Rasio and Shapiro [26] may be used to check the validity of numerical solutions. However, in numerical solutions of Uryū and Eriguchi the velocity component along the orbital axis exists while in those of Lai, Rasio and Shapiro such a component is assumed to be zero from the beginning. Therefore new analytic or semi-analytic solutions are needed to check numerical solutions even in

Newtonian gravity. Such a check of numerical solutions is extremely important because in the numerical calculation, there is a possibility to obtain another solution although the binding energy of a binary neutron star is almost the same value, and to lead a different conclusion [27,28].

In the previous paper [29], we showed such a new, almost analytic solution to an equilibrium of irrotational binary polytropic stars for the polytropic index $n = 1$ in Newtonian gravity by expanding all physical quantities in a power of $\epsilon \equiv a_0/R$, where R and a_0 are the orbital separation of the binary system and the radius of each star for $R = \infty$. In that paper, we have extended the method developed by Chandrasekhar more than 65 years ago for corotating fluids [30,31] to the one for irrotational fluids. In this paper, we show semi-analytic solutions for arbitrary polytropic indices by numerically solving ordinary differential equations.

This paper is organized as follows. In §2, we formulate the method to solve the irrotational binary polytropic stars. In §3, the physical values, i.e., the central density of a star, the orbital angular velocity, the total energy and total angular momentum of the binary system are calculated. The numerical results are presented in §4. Section 5 is devoted to summary and discussions.

Although a binary system consists of two stars, we pay particular attention to one of two stars. We call it star 1 whose mass is M_1 and the companion one star 2 whose mass is M_2 . In this paper, we adopt two corotating coordinate systems. First one is a Cartesian coordinate system $\mathbf{X}$ whose origin is located at the center of mass of the binary system. For calculational convenience, we choose the orbital axis as X_3 , and we take the direction of X_1 from the center of mass of star 2 to that of star 1. The second coordinate system is the spherical one $\mathbf{r} = (r, \theta, \varphi)$ whose origin is located at the center of mass of star 1. We use units of $G = 1$.

II. FORMULATION

Since we treat irrotational fluids in Newtonian gravity, the basic equations for obtaining equilibrium configurations of binary systems are the equation of state, the Euler equation, the equation of continuity and the Poisson equation:

$$P = K\rho^{1+\frac{1}{n}}, \quad (2.1)$$

$$\nabla \left[\int \frac{dP}{\rho} - U + \frac{1}{2}v^2 - \mathbf{v} \cdot (\boldsymbol{\Omega} \times \mathbf{r}) \right] = 0, \quad (2.2)$$

$$\nabla \cdot \mathbf{v} = -(\mathbf{v} - \boldsymbol{\Omega} \times \mathbf{r}) \cdot \frac{\nabla \rho}{\rho}, \quad (2.3)$$

$$\Delta U = -4\pi\rho, \quad (2.4)$$

where P , ρ , n , U and Ω are the pressure, the density, the polytropic index, the gravitational potential and the orbital angular velocity, respectively. K is a constant related to entropy and $\mathbf{v}$ represents the velocity field in the inertial frame. The gravitational potential U is separated into two parts, i.e., the contribution from star 1 to itself $U^{1 \rightarrow 1}$ and that from star 2 to star 1 $U^{2 \rightarrow 1}$. We can also express the gravitational potential contributed from star 1 to star 2 as $U^{1 \rightarrow 2}$. These gravitational potentials are written as

$$U^{2 \rightarrow 1} = \frac{M_2}{R} \sum_{l=0}^{\infty} (-1)^l \left(\frac{r}{R} \right)^l P_l(\sin \theta \cos \varphi) + \frac{3\mathcal{I}_{11}'}{2R^3} \left[1 - 3 \left(\frac{r}{R} \right) P_1(\sin \theta \cos \varphi) + O(R^{-2}) \right] + \text{higher order terms}, \quad (2.5)$$

$$U^{1 \rightarrow 2} = \frac{M_1}{R} \sum_{l=0}^{\infty} \left(\frac{r'}{R} \right)^l P_l(\sin \theta' \cos \varphi') + \frac{3\mathcal{I}_{11}}{2R^3} \left[1 + 3 \left(\frac{r'}{R} \right) P_1(\sin \theta' \cos \varphi') + O(R^{-2}) \right] + \text{higher order terms}, \quad (2.6)$$

where the superscript $'$ means the term concerned with star 2 and P_l denotes the Legendre function of order l . $\mathcal{I}_{11}$ is the reduced quadrupole moment defined by

$$\begin{aligned} \mathcal{I}_{11} &= \frac{1}{3} (2I_{11} - I_{22} - I_{33}), \\ &= \frac{2}{3} \int_{\text{star 1}} d^3x \rho r^2 P_2(\sin \theta \cos \varphi). \end{aligned} \quad (2.7)$$

In the irrotational fluid case, we can express $\mathbf{v}$ as a gradient of a scalar function Φ , i.e.,

$$\mathbf{v} = \nabla\Phi. \quad (2.8)$$

Following the *Lane-Emden equation* we first express the density as

$$\rho = \rho_c \Theta^n(\xi, \theta, \varphi) \quad (2.9)$$

with $\xi = r/\alpha$. Here the definitions of α and α' are

$$\alpha \equiv \left[\frac{K(1+n)\rho_c^{\frac{1}{n}-1}}{4\pi} \right]^{1/2}, \quad (2.10)$$

$$\alpha' \equiv \left[\frac{K'(1+n')\rho_c'^{\frac{1}{n'}-1}}{4\pi} \right]^{1/2}, \quad (2.11)$$

with ρ_c being the central density. Let us define $\epsilon \equiv a_0/R$, where R and a_0 are the orbital separation of the binary system and the radius of star 1 for $R = \infty$ [29]. The radius of a spherical star a_0 is given by $a_0 = \alpha\xi_1$ as usual. Also we define $\epsilon' \equiv a'_0/R$ for star 2. Note that in the case of Newtonian gravity, we have two degrees of freedom for describing the differences of mass, i.e., ρ_c and α . Therefore, we can express the differences of mass by changing only ρ_c with fixing α . In this case, we can make the radius of each star and each expansion parameter coincide. Although we have adopted such a situation in Ref. [29], in the present paper we express equations concerned with star 2 by using ϵ' , α' and so on.

We expand Θ in a power series of a parameter ϵ as¹

$$\Theta = \sum_{i=0}^{\infty} \epsilon^i \Theta_i. \quad (2.12)$$

Since the shape of star 1 is spherical when R is large, the lowest order term of Θ is the solution of the *Lane-Emden equation*. Then we expand Θ_i by spherical harmonics as

$$\Theta_i = \sum_{l,m} {}^{(i)}\psi_{lm}(\xi) Y_l^m(\theta, \varphi). \quad (2.13)$$

Now we consider the orbital motion of star 1. In the spherical coordinate system, it becomes

$$\boldsymbol{\Omega} \times \mathbf{r} = \Omega R (\tilde{\boldsymbol{\Omega}} \times \boldsymbol{\xi})_{orb} + \Omega \alpha (\tilde{\boldsymbol{\Omega}} \times \boldsymbol{\xi})_{fig}, \quad (2.14)$$

where

$$(\tilde{\boldsymbol{\Omega}} \times \boldsymbol{\xi})_{orb} = \frac{1}{1+p} (\sin \theta \sin \varphi, \cos \theta \sin \varphi, \cos \varphi), \quad (2.15)$$

$$(\tilde{\boldsymbol{\Omega}} \times \boldsymbol{\xi})_{fig} = (0, 0, \xi \sin \theta), \quad (2.16)$$

and $p \equiv M_1/M_2$. The first term on the right-hand side of Eq. (2.14) comes from the orbital motion of the center of mass of star 1 and the second term comes from the fluid motion around the center of mass of star 1. The orbital motion of star 2 becomes

$$\boldsymbol{\Omega} \times \mathbf{r}' = \Omega R (\tilde{\boldsymbol{\Omega}} \times \boldsymbol{\xi})'_{orb} + \Omega \alpha' (\tilde{\boldsymbol{\Omega}} \times \boldsymbol{\xi})'_{fig}, \quad (2.17)$$

where

$$(\tilde{\boldsymbol{\Omega}} \times \boldsymbol{\xi})'_{orb} = -\frac{p}{1+p} (\sin \theta' \sin \varphi', \cos \theta' \sin \varphi', \cos \varphi'), \quad (2.18)$$

$$(\tilde{\boldsymbol{\Omega}} \times \boldsymbol{\xi})'_{fig} = (0, 0, \xi' \sin \theta'), \quad (2.19)$$

¹Note that we express the Lane-Emden function for star 2 as $\bar{\Theta}$ and expand it in a power series of a parameter ϵ' as $\bar{\Theta} = \sum_i \epsilon'^i \bar{\Theta}_i$.

in the spherical coordinate system whose origin is located at the center of mass of star 2. Here we have expressed the spherical coordinate system for star 2 as $\mathbf{r}' = (r', \theta', \varphi')$ with $\xi' = r'/\alpha'$.

Next, we rewrite the equation of continuity (2.3) as

$$\Delta\Phi = -n(\nabla\Phi - \boldsymbol{\Omega} \times \mathbf{r}) \cdot \frac{\nabla\Theta}{\Theta}. \quad (2.20)$$

The condition for Φ at the stellar surface is

$$(\nabla\Phi - \boldsymbol{\Omega} \times \mathbf{r}) \cdot (\nabla\Theta)|_{surf} = 0, \quad (2.21)$$

since $\Theta = 0$ at the surface. We expand Φ also as

$$\Phi = \sum_{i=0}^{\infty} \epsilon^i \Phi_i. \quad (2.22)$$

When R is large, the shape of star 1 is spherical and star 1 has only the orbital motion of the center of mass in the inertial frame with no intrinsic spin so that the gradient of the lowest order term of Φ should agree with the orbital motion of the center of mass of star 1, i.e.,

$$\nabla\Phi_0 = \Omega R(\tilde{\boldsymbol{\Omega}} \times \boldsymbol{\xi})_{orb}. \quad (2.23)$$

Considering Φ_0 , we normalize Φ as

$$\tilde{\Phi} = \frac{\epsilon\Phi}{\Omega\alpha a_0}. \quad (2.24)$$

We again expand $\tilde{\Phi}_i$ by spherical harmonics as

$$\tilde{\Phi}_i = \sum_{l,m} {}^{(i)}\phi_{lm}(\xi) \tilde{Y}_l^m(\theta, \varphi), \quad (2.25)$$

where $\tilde{Y}_l^m$ is defined by using the spherical harmonics Y_l^m as

$$\tilde{Y}_l^m(\theta, \varphi) \equiv C_{lm} \frac{\partial}{\partial \varphi} Y_l^m(\theta, \varphi) \quad (2.26)$$

with constants C_{lm} . Note that the gradient of the lowest order term of Φ has order ϵ^{-1} except for the implicit dependence in Ω . When we normalize Φ using Eq. (2.24), the lowest order term of $\tilde{\Phi}$ has order ϵ^0 . We must keep this difference in mind.

The orbital angular velocity is derived from the force balance equation, i.e., the first tensor virial relation, defined by [32]

$$\int_{star\ 1} d^3x \frac{\partial P}{\partial x_1} = 0, \quad (2.27)$$

where $x_1 = r \sin \theta \cos \varphi$. This equation is expressed by using Eq. (2.2) as

$$0 = \int_{star\ 1} d^3x \rho \left[\frac{\partial U^{1 \rightarrow 1}}{\partial x_1} + \frac{\partial U^{2 \rightarrow 1}}{\partial x_1} - \frac{1}{2} \frac{\partial}{\partial x_1} (\nabla\Phi)^2 + \frac{\partial}{\partial x_1} \left\{ (\nabla\Phi) \cdot (\boldsymbol{\Omega} \times \mathbf{r}) \right\} \right]. \quad (2.28)$$

If we substitute Eq. (2.14), Θ_0 and Φ_0 into Eq. (2.28), we obtain the orbital angular velocity in the lowest order as

$$\Omega_0^2 = \frac{M_{tot}}{R^3}, \quad (2.29)$$

where $M_{tot} = M_1 + M_2$. Note here that we also expand Ω^2 as

$$\Omega^2 = \sum_{i=0}^{\infty} \epsilon^i \Omega_i^2. \quad (2.30)$$

Finally, we express the gravitational potential by rewriting Eq. (2.2) as

$$U = K(1+n)\rho^{\frac{1}{n}} + \frac{1}{2}v^2 - \mathbf{v} \cdot (\mathbf{\Omega} \times \mathbf{r}) + U_0, \quad (2.31)$$

where U_0 is constant. Substituting Eq. (2.31) into the Poisson equation (2.4), we obtain the equation to determine the equilibrium figure as

$$\alpha^2 \Delta \Theta = -\Theta^n - \frac{1}{8\pi\rho_c} \Delta \left[(\nabla \Phi)^2 - 2(\nabla \Phi) \cdot (\mathbf{\Omega} \times \mathbf{r}) \right]. \quad (2.32)$$

Now a solution can be obtained iteratively. Firstly, Θ_i is determined by demanding that the gravitational potential and its normal derivative are continuous at the stellar surface [30], that is,

$$U_{int}(\xi = \Xi) = U_{ext}(\xi = \Xi), \quad (2.33)$$

$$\frac{\partial U_{int}}{\partial \xi}(\xi = \Xi) = \frac{\partial U_{ext}}{\partial \xi}(\xi = \Xi), \quad (2.34)$$

where $\Xi(\theta, \varphi)$ expresses the surface,

$$\Theta(\xi = \Xi(\theta, \varphi)) = 0. \quad (2.35)$$

Substituting Θ_i and Eq. (2.14) into Eqs. (2.20) and (2.28), we obtain Φ_i and Ω_i^2 . Secondly we substitute Φ_i and Ω_i^2 into Eq. (2.32) and derive Θ_{i+1} . We have continued this procedure up to order ϵ^6 in this paper.

III. DERIVATION OF THE PHYSICAL VALUES

In this section, we calculate physical values such as the baryon mass of star 1, the orbital angular velocity, the total energy and total angular momentum of the binary system. We will just show the main results. All the details will be shown in the Appendix.

A. Mass and Central Density

The baryon mass of star 1 is calculated by integrating the density over the volume of star 1 as

$$M_1 = \int_{star\ 1} d^3x \rho, \quad (3.1)$$

$$\begin{aligned} &= \alpha^3 \rho_c \int d\xi d\theta d\varphi \xi^2 \sin \theta \left[\Theta_0^n + \epsilon^3 n \Theta_0^{n-1} \Theta_3 + \epsilon^4 n \Theta_0^{n-1} \Theta_4 + \epsilon^5 n \Theta_0^{n-1} \Theta_5 \right. \\ &\quad \left. + \epsilon^6 n \left\{ \Theta_0^{n-1} \Theta_6 + \frac{1}{2}(n-1) \Theta_0^{n-2} \Theta_3^2 \right\} \right]. \end{aligned} \quad (3.2)$$

Note here that the integration range for ξ is $[0, \Xi(\theta, \varphi)]$.

For $n > 0$ case, the mass of star 1 becomes

$$M_1 = -4\pi\rho_c\alpha^3\xi_1^2\Theta_{0,\xi}(\xi_1) \left[1 + \epsilon^6 \left(\frac{1}{\Theta_{0,\xi}(\xi_1)} \right) \left(\frac{({}^{(3)}\psi_2(\xi_1))^2}{10\Theta_{0,\xi}(\xi_1)} n \Theta_0^{n-1}(\xi_1) + {}^{(6)}\psi_{0,\xi}(\xi_1) \right) \right], \quad (3.3)$$

$$= -4\pi\rho_c\alpha^3\xi_1^2\Theta_{0,\xi}(\xi_1) \left[1 + \epsilon^6 \left(\frac{-1}{\xi_1\Theta_{0,\xi}(\xi_1)} \right) \left(\tilde{c}_6 + {}^{(6)}\psi_0(\xi_1) \right) \right]. \quad (3.4)$$

where we define

$$_{,\xi} \equiv \frac{d}{d\xi}. \quad (3.5)$$

We have used Eq. (H43) in deriving Eq. (3.4) from Eq. (3.3). Here $\tilde{c}_6$ is a constant, and one can see the expression of $\tilde{c}_6$ in Appendix H. We assume that the baryon mass $M_1 = -4\pi\rho_{c0}\alpha_0^3\xi_1^2\Theta_{0,\xi}(\xi_1)$ is conserved throughout the

equilibrium sequences of the binary system, where ρ_{c0} and α_0 denote the values for a spherical star. Therefore, the central density and α are expressed as

$$\rho_c = \rho_{c0} \left(1 - \epsilon^6 \frac{\delta \rho_c}{\rho_c} \right), \quad (3.6)$$

$$\alpha = \left[\frac{K(1+n)\rho_{c0}^{\frac{1}{n}-1}}{4\pi} \right]^{1/2} \left(1 - \epsilon^6 \frac{1-n}{2n} \frac{\delta \rho_c}{\rho_c} \right), \quad (3.7)$$

where we define²

$$\frac{\delta \rho_c}{\rho_c} \equiv \frac{2n}{3-n} \left(\frac{-1}{\xi_1 \Theta_{0,\xi}(\xi_1)} \right) \left(\tilde{c}_6 + {}^{(6)}\psi_0(\xi_1) \right). \quad (3.8)$$

B. Orbital angular velocity

We obtain the orbital angular velocity from Eq. (2.28) as

$$\begin{aligned} \Omega^2 &= \frac{M_{tot}}{R^3} \left[1 + \frac{9}{2R^2} \left(\epsilon^3 \frac{\bar{\mathcal{I}}_{11}}{M_1} + \epsilon'^3 \frac{\bar{\mathcal{I}}'_{11}}{M_2} \right) + O(\epsilon^7) \right], \\ &= \frac{M_{tot}}{a_0^3} \epsilon^3 \left[1 + \frac{9\epsilon^2}{2a_0^2} \left(\epsilon^3 \frac{\bar{\mathcal{I}}_{11}}{M_1} + \epsilon'^3 \frac{\bar{\mathcal{I}}'_{11}}{M_2} \right) + O(\epsilon^7) \right], \end{aligned} \quad (3.9)$$

where $\bar{\mathcal{I}}_{11}$ denotes $\mathcal{I}_{11}/\epsilon^3$.

C. Total energy

The total energy is written as

$$E = \Pi_{tot} + (W_{self})_{tot} + (W_{int})_{tot} + T_{tot}, \quad (3.10)$$

where Π_{tot} , $(W_{self})_{tot}$, $(W_{int})_{tot}$ and T_{tot} denote the total internal energy, the total self-gravity energy, the total interaction energy and the total kinetic energy, respectively. We calculate each energy in the following.

1. Internal energy

The definition of the internal energy of star 1 is

$$\Pi_1 = n \int_{star\ 1} d^3x P. \quad (3.11)$$

For $n > 0$ case, we obtain

$$\Pi_1 = \frac{n}{1+n} \frac{M_1^2}{a_0 \xi_1^3 (\Theta_{0,\xi}(\xi_1))^2} \left[\left(1 - \epsilon^6 \frac{5-n}{2n} \frac{\delta \rho_c}{\rho_c} \right) \int_0^{\xi_1} d\xi \xi^2 \Theta_0^{1+n} + \epsilon^6 (1+n) \int_0^{\xi_1} d\xi \xi^2 \left\{ \Theta_0^{n(6)} \psi_0 + \frac{1}{10} n \Theta_0^{n-1} ({}^{(3)}\psi_2)^2 \right\} \right]. \quad (3.12)$$

Therefore, the total internal energy is expressed as

²The expression of $\delta \rho_c / \rho_c$ seems to diverge if $n < 1$ when one see Eq. (3.3). However physically $\delta \rho_c / \rho_c$ should be finite. In reality we determined $\delta \rho_c / \rho_c$ by using the virial relation as shown in the Appendix I.

$$\begin{aligned}\Pi_{tot} = & \frac{n}{1+n} \frac{M_1^2}{a_0 \xi_1^3 (\Theta_{0,\xi}(\xi_1))^2} \left[\left(1 - \epsilon^6 \frac{5-n}{2n} \frac{\delta \rho_c}{\rho_c} \right) \int_0^{\xi_1} d\xi \xi^2 \Theta_0^{1+n} + \epsilon^6 (1+n) \int_0^{\xi_1} d\xi \xi^2 \left\{ \Theta_0^{n(6)} \psi_0 + \frac{1}{10} n \Theta_0^{n-1} ((^{(3)}\psi_2)^2) \right\} \right] \\ & + \frac{n'}{1+n'} \frac{M_2^2}{a'_0 \xi_1'^3 (\bar{\Theta}_{0,\xi'}(\xi_1'))^2} \left[\left(1 - \epsilon'^6 \frac{5-n'}{2n'} \frac{\delta \rho'_c}{\rho'_c} \right) \int_0^{\xi_1'} d\xi' \xi'^2 \bar{\Theta}_0^{1+n'} \right. \\ & \left. + \epsilon'^6 (1+n') \int_0^{\xi_1'} d\xi' \xi'^2 \left\{ \bar{\Theta}_0^{n'(6)} \bar{\psi}_0 + \frac{1}{10} n' \bar{\Theta}_0^{n'-1} ((^{(3)}\bar{\psi}_2)^2) \right\} \right],\end{aligned}\tag{3.13}$$

where the superscript ' means the terms concerned with star 2, and $\bar{}$ also means the functions concerned with star 2.

2. Self-gravity energy

The definition of the self-gravity energy of star 1 is

$$W_{self,1} = -\frac{1}{2} \int_{star\ 1} d^3x \rho U^{1 \rightarrow 1}.\tag{3.14}$$

For $n > 0$ case, we obtain

$$W_{self,1} = -\frac{1+n}{2n} \Pi_1 + \epsilon^6 \pi \rho_c \left[\frac{3\mu \bar{F}_{11}}{\xi_1^2 (1+p)} - 2\alpha^2 M_1 \tilde{c}_6 \right] + 2\pi \rho_c \alpha^2 M_1 \xi_1 \Theta_{0,\xi}(\xi_1),\tag{3.15}$$

where μ is defined as

$$\mu \equiv \frac{M_{tot}}{4\pi \rho_c \alpha^3 \xi_1} = -\left(\frac{1+p}{p} \right) \xi_1 \Theta_{0,\xi}(\xi_1).\tag{3.16}$$

The self-gravity energy of star 2 is

$$W_{self,2} = -\frac{1+n'}{2n'} \Pi_2 + \epsilon'^6 \pi \rho'_c \left[\frac{3\mu' p \bar{F}'_{11}}{\xi_1'^2 (1+p)} - 2\alpha'^2 M_2 \tilde{c}'_6 \right] + 2\pi \rho'_c \alpha'^2 M_2 \xi_1' \bar{\Theta}_{0,\xi'}(\xi_1').\tag{3.17}$$

Here, $\tilde{c}'_6$ is a constant similar to $\tilde{c}_6$, and

$$\mu' \equiv \frac{M_{tot}}{4\pi \rho'_c \alpha'^3 \xi_1'} = -(1+p) \xi_1' \bar{\Theta}_{0,\xi'}(\xi_1').\tag{3.18}$$

Therefore, the total self-gravity energy becomes

$$\begin{aligned}(W_{self})_{tot} = & -\frac{1+n}{2n} \Pi_1 + \epsilon^6 \frac{M_1^2}{2a_0} \left(\frac{3}{2p} \frac{\bar{F}_{11}}{M_1 a_0^2} + \frac{\tilde{c}_6}{\xi_1 \Theta_{0,\xi}(\xi_1)} \right) - \frac{M_1^2}{2a_0} \left(1 - \epsilon^6 \frac{1}{n} \frac{\delta \rho_c}{\rho_c} \right) \\ & - \frac{1+n'}{2n'} \Pi_2 + \epsilon'^6 \frac{M_2^2}{2a_0'} \left(\frac{3p}{2} \frac{\bar{F}'_{11}}{M_2 a_0'^2} + \frac{\tilde{c}'_6}{\xi_1' \bar{\Theta}_{0,\xi'}(\xi_1')} \right) - \frac{M_2^2}{2a_0'} \left(1 - \epsilon'^6 \frac{1}{n'} \frac{\delta \rho'_c}{\rho'_c} \right).\end{aligned}\tag{3.19}$$

3. Interaction energy

The definition of the interaction energy of star 1 is

$$W_{int,1} = -\frac{1}{2} \int_{star\ 1} d^3x \rho U^{2 \rightarrow 1},\tag{3.20}$$

$$= -\frac{M_1 M_2}{2a_0} \epsilon - \epsilon^3 \frac{3M_1 M_2}{4a_0^3} \left(\epsilon^3 \frac{\bar{F}_{11}}{M_1} + \epsilon'^3 \frac{\bar{F}'_{11}}{M_2} \right).\tag{3.21}$$

Therefore, the total interaction energy becomes

$$(W_{int})_{tot} = -\frac{M_1 M_2}{a_0} \epsilon - \epsilon^3 \frac{3M_1 M_2}{2a_0^3} \left(\epsilon^3 \frac{\bar{F}_{11}}{M_1} + \epsilon'^3 \frac{\bar{F}'_{11}}{M_2} \right).\tag{3.22}$$

4. Kinetic energy

The definition of the kinetic energy of star 1 is

$$T_1 = \frac{1}{2} \int_{star\ 1} d^3x \rho v_1^2, \quad (3.23)$$

$$= \frac{M_1 M_2}{2a_0(1+p)} \epsilon \left[1 + \frac{9\epsilon^2}{2a_0^2} \left(\epsilon^3 \frac{\bar{F}_{11}}{M_1} + \epsilon'^3 \frac{\bar{F}'_{11}}{M_2} \right) \right]. \quad (3.24)$$

The kinetic energy of star 2 is also given as

$$T_2 = \frac{M_1 M_2 p}{2a_0(1+p)} \epsilon \left[1 + \frac{9\epsilon^2}{2a_0^2} \left(\epsilon^3 \frac{\bar{F}_{11}}{M_1} + \epsilon'^3 \frac{\bar{F}'_{11}}{M_2} \right) \right]. \quad (3.25)$$

Then the total kinetic energy becomes

$$T_{tot} = \frac{M_1 M_2}{2a_0} \epsilon + \epsilon^3 \frac{9M_1 M_2}{4a_0^3} \left(\epsilon^3 \frac{\bar{F}_{11}}{M_1} + \epsilon'^3 \frac{\bar{F}'_{11}}{M_2} \right). \quad (3.26)$$

D. Total angular momentum

The total angular momentum is calculated from the equation;

$$\mathbf{J} = \int_{star\ 1, star\ 2} d^3x \rho (\mathbf{R} \times \mathbf{v}), \quad (3.27)$$

where $\mathbf{R}$ for star 1 is written as

$$\mathbf{R} = \frac{R}{1+p} (\sin \theta \cos \varphi, \cos \theta \cos \varphi, -\sin \varphi) + r(1, 0, 0), \quad (3.28)$$

and for star 2 is

$$\mathbf{R}' = -\frac{pR}{1+p} (\sin \theta' \cos \varphi', \cos \theta' \cos \varphi', -\sin \varphi') + r'(1, 0, 0) \quad (3.29)$$

in the spherical coordinate system. Note that since the velocity field $\mathbf{v} = \nabla \Phi$ is written in the spherical coordinate system, it is convenient to calculate $\mathbf{R} \times \mathbf{v}$ in the spherical one. However, we should be careful to integrate it over the volume of each star, because in the spherical coordinate system, the unit vectors depend on the coordinates. For example, the unit vector $(1, 0, 0)$ in the *Cartesian* coordinate system becomes $(\sin \theta \cos \varphi, \cos \theta \cos \varphi, -\sin \varphi)$ in the *spherical* one.

As a result, we obtain the total angular momentum;

$$\mathbf{J} = (0, 0, J) \quad (3.30)$$

in the *Cartesian* coordinate system. Here J is calculated as

$$J = \frac{M_1 M_2}{M_1 + M_2} R^2 \Omega \left[1 + \text{higher order terms than } O(\epsilon^6) \right]. \quad (3.31)$$

E. Virial Relation

By using the above energies, we can calculate the virial relation which is written as

$$\frac{3}{n} \Pi_1 + \frac{3}{n'} \Pi_2 + (W_{self})_{tot} + (W_{int})_{tot} + 2T_{tot} = 0. \quad (3.32)$$

This equation is used to check the solutions as shown in the previous letter [29]. However in this paper we use it to determine $\delta\rho_c/\rho_c$ (see Appendix I).

IV. NUMERICAL RESULTS

In the following sections, we set $M_1 = M_2 = M$, $n = n'$, $\alpha = \alpha'$, $\rho_c = \rho'_c$ and so on, i.e., the identical star binary for simplicity, and investigate the cases of different polytropic indices.

First, we show the Lane-Emden function of star 1 up to $O(\epsilon^6)$. It is written as

$$\Theta = \Theta_0 + \epsilon^3 \Theta_3 + \epsilon^4 \Theta_4 + \epsilon^5 \Theta_5 + \epsilon^6 \Theta_6, \quad (4.1)$$

where

$$\Theta_3 = {}^{(3)}\psi_2(\xi) P_2(\sin \theta \cos \varphi), \quad (4.2)$$

$$\Theta_4 = {}^{(4)}\psi_3(\xi) P_3(\sin \theta \cos \varphi), \quad (4.3)$$

$$\Theta_5 = {}^{(5)}\psi_4(\xi) P_4(\sin \theta \cos \varphi), \quad (4.4)$$

$$\begin{aligned} \Theta_6 = & {}^{(6)}\psi_0(\xi) + {}^{(6)}\psi_2(\xi) P_2(\sin \theta \cos \varphi) + {}^{(6)}\psi_{22}(\xi) P_2^2(\cos \theta) \cos 2\varphi \\ & + {}^{(6)}\psi_4(\xi) P_4(\sin \theta \cos \varphi) + {}^{(6)}\psi_5(\xi) P_5(\sin \theta \cos \varphi). \end{aligned} \quad (4.5)$$

The functions Θ_0 , ${}^{(i)}\psi_j$ and their derivatives for each polytropic index are given in Tables I–VIII.

Next, we give the velocity potential up to $O(\epsilon^6)$. We can write it as

$$\tilde{\Phi} = \tilde{\Phi}_0 + \epsilon^4 \tilde{\Phi}_4 + \epsilon^5 \tilde{\Phi}_5 + \epsilon^6 \tilde{\Phi}_6, \quad (4.6)$$

where

$$\tilde{\Phi}_0 = \frac{1}{1+p} \xi \sin \theta \sin \varphi, \quad (4.7)$$

$$\tilde{\Phi}_4 = {}^{(4)}\phi_2(\xi) P_2^2(\cos \theta) \sin 2\varphi, \quad (4.8)$$

$$\tilde{\Phi}_5 = {}^{(5)}\phi_3(\xi) \left[P_3^1(\cos \theta) \sin \varphi - \frac{1}{2} P_3^3(\cos \theta) \sin 3\varphi \right], \quad (4.9)$$

$$\tilde{\Phi}_6 = {}^{(6)}\phi_4(\xi) \left[P_4^2(\cos \theta) \sin 2\varphi - \frac{1}{4} P_4^4(\cos \theta) \sin 4\varphi \right]. \quad (4.10)$$

The functions ${}^{(4)}\phi_2$, ${}^{(5)}\phi_3$ and ${}^{(6)}\phi_4$ and their derivatives are shown in Tables IX–XII. The derivatives are related to the velocity field in the inertial frame. Note here that in Eq. (4.7) we write the dependence of the mass ratio $p = M_1/M_2$ although we consider only $p = 1$ case in this section.

Here we directly show the existence of the velocity component along the orbital axis. In this paper, we take X_3 as the orbital axis. Since the origin of the corotating coordinate system $\mathbf{X}$ is located at the center of mass of the binary system, it is convenient to consider another Cartesian coordinate system which origin is located at the center of mass of star 1. We call it $\mathbf{x} = (x_1, x_2, x_3)$. The relations between $\mathbf{x}$ and the spherical coordinate system $\mathbf{r}$ are

$$x_1 = r \sin \theta \cos \varphi, \quad (4.11)$$

$$x_2 = r \sin \theta \sin \varphi, \quad (4.12)$$

$$x_3 = r \cos \theta, \quad (4.13)$$

as usual. Then, the velocity component along the orbital axis in the inertial frame is written as

$$v_3 = \frac{\partial \Phi}{\partial x_3}. \quad (4.14)$$

In the corotating frame, we can express the velocity field $\mathbf{u}$ as

$$\mathbf{u} = \nabla \Phi - \boldsymbol{\Omega} \times \mathbf{r}, \quad (4.15)$$

$$= -\Omega \alpha (\tilde{\boldsymbol{\Omega}} \times \boldsymbol{\xi})_{fig} + \epsilon^4 \nabla \Phi_4 + \epsilon^5 \nabla \Phi_5 + \epsilon^6 \nabla \Phi_6, \quad (4.16)$$

$$= -\Omega \alpha (\tilde{\boldsymbol{\Omega}} \times \boldsymbol{\xi})_{fig} + \Omega a_0 \left[\epsilon^3 \tilde{\nabla} \tilde{\Phi}_4 + \epsilon^4 \tilde{\nabla} \tilde{\Phi}_5 + \epsilon^5 \tilde{\nabla} \tilde{\Phi}_6 \right], \quad (4.17)$$

where $\tilde{\nabla} = \alpha \nabla$ is a nabla defined by using (ξ, θ, φ) and given in Eq. (A4). Then, the velocity component along the orbital axis in the corotating frame becomes

$$u_3 = \Omega a_0 \left[\epsilon^3 \frac{\partial \tilde{\Phi}_4}{\partial \tilde{x}_3} + \epsilon^4 \frac{\partial \tilde{\Phi}_5}{\partial \tilde{x}_3} + \epsilon^5 \frac{\partial \tilde{\Phi}_6}{\partial \tilde{x}_3} \right], \quad (4.18)$$

where we have used the normalized coordinate system,

$$\tilde{x}_1 = \xi \sin \theta \cos \varphi, \quad (4.19)$$

$$\tilde{x}_2 = \xi \sin \theta \sin \varphi, \quad (4.20)$$

$$\tilde{x}_3 = \xi \cos \theta, \quad (4.21)$$

for the calculational convenience. Note that since $\nabla \Phi$ is proportional to Ω/ϵ , the equation (4.16) does not express the real dependence on ϵ .

Next, we calculate the term for $\tilde{\Phi}_4$ in Eq. (4.18). Then we obtain

$$\begin{aligned} \frac{\partial \tilde{\Phi}_4}{\partial \tilde{x}_3} &= \cos \theta \frac{\partial \tilde{\Phi}_4}{\partial \xi} - \frac{\sin \theta}{\xi} \frac{\partial \tilde{\Phi}_4}{\partial \theta}, \\ &= 3 \left(\frac{d^{(4)}\phi_2}{d\xi} - \frac{2^{(4)}\phi_2}{\xi} \right) \sin^2 \theta \cos \theta \sin 2\varphi. \end{aligned} \quad (4.22)$$

If $^{(4)}\phi_2(\xi)$ is proportional to ξ^2 , the velocity component along the orbital axis disappears³. However, $^{(4)}\phi_2(\xi)$ is not proportional to ξ^2 at all for $n > 0$. Moreover, we can show the velocity components along the orbital axis for Φ_5 and Φ_6 as

$$\frac{\partial \tilde{\Phi}_5}{\partial \tilde{x}_3} = 6 \left[\left(\frac{d^{(5)}\phi_3}{d\xi} - \frac{3^{(5)}\phi_3}{\xi} \right) (1 - 5 \sin^2 \theta \cos^2 \varphi) + \frac{2^{(5)}\phi_3}{\xi} \right] \sin \theta \cos \theta \sin \varphi, \quad (4.23)$$

$$\frac{\partial \tilde{\Phi}_6}{\partial \tilde{x}_3} = 30 \left[\left(\frac{d^{(6)}\phi_4}{d\xi} - \frac{4^{(6)}\phi_4}{\xi} \right) (3 - 7 \sin^2 \theta \cos^2 \varphi) + \frac{6^{(6)}\phi_4}{\xi} \right] \sin^2 \theta \cos \theta \sin \varphi \cos \varphi. \quad (4.24)$$

In these equations, we can see that even if $^{(5)}\phi_3 \propto \xi^3$ and $^{(6)}\phi_4 \propto \xi^4$, there remain the velocity components along the orbital axis. The velocity component along the orbital axis is order ϵ^3 higher. However, if one would like to pay attention to the internal state of neutron stars in the binary system, it should be taken into account.

We give also the other components of the velocity field as

$$u_1 = \Omega x_2 + \Omega a_0 \left[\epsilon^3 \frac{\partial \tilde{\Phi}_4}{\partial \tilde{x}_1} + \epsilon^4 \frac{\partial \tilde{\Phi}_5}{\partial \tilde{x}_1} + \epsilon^5 \frac{\partial \tilde{\Phi}_6}{\partial \tilde{x}_1} \right], \quad (4.25)$$

$$u_2 = -\Omega x_1 + \Omega a_0 \left[\epsilon^3 \frac{\partial \tilde{\Phi}_4}{\partial \tilde{x}_2} + \epsilon^4 \frac{\partial \tilde{\Phi}_5}{\partial \tilde{x}_2} + \epsilon^5 \frac{\partial \tilde{\Phi}_6}{\partial \tilde{x}_2} \right], \quad (4.26)$$

where

$$\frac{\partial \tilde{\Phi}_4}{\partial \tilde{x}_1} = \sin \theta \cos \varphi \frac{\partial \tilde{\Phi}_4}{\partial \xi} + \frac{\cos \theta \cos \varphi}{\xi} \frac{\partial \tilde{\Phi}_4}{\partial \theta} - \frac{\sin \varphi}{\xi \sin \theta} \frac{\partial \tilde{\Phi}_4}{\partial \varphi}, \quad (4.27)$$

$$= 6 \left[\left(\frac{d^{(4)}\phi_2}{d\xi} - \frac{2^{(4)}\phi_2}{\xi} \right) \sin^2 \theta \cos^2 \varphi + \frac{^{(4)}\phi_2}{\xi} \right] \sin \theta \sin \varphi, \quad (4.28)$$

$$\frac{\partial \tilde{\Phi}_4}{\partial \tilde{x}_2} = \sin \theta \sin \varphi \frac{\partial \tilde{\Phi}_4}{\partial \xi} + \frac{\cos \theta \sin \varphi}{\xi} \frac{\partial \tilde{\Phi}_4}{\partial \theta} + \frac{\cos \varphi}{\xi \sin \theta} \frac{\partial \tilde{\Phi}_4}{\partial \varphi}, \quad (4.29)$$

$$= 6 \left[\left(\frac{d^{(4)}\phi_2}{d\xi} - \frac{2^{(4)}\phi_2}{\xi} \right) \sin^2 \theta \sin^2 \varphi + \frac{^{(4)}\phi_2}{\xi} \right] \sin \theta \cos \varphi, \quad (4.30)$$

$$\frac{\partial \tilde{\Phi}_5}{\partial \tilde{x}_1} = 6 \left[\left(\frac{d^{(5)}\phi_3}{d\xi} - \frac{3^{(5)}\phi_3}{\xi} \right) (1 - 5 \sin^2 \theta \cos^2 \varphi) - \frac{8^{(5)}\phi_3}{\xi} \right] \sin^2 \theta \sin \varphi \cos \varphi, \quad (4.31)$$

$$\frac{\partial \tilde{\Phi}_5}{\partial \tilde{x}_2} = 6 \left[\left(\frac{d^{(5)}\phi_3}{d\xi} - \frac{3^{(5)}\phi_3}{\xi} \right) (1 - 5 \sin^2 \theta \cos^2 \varphi) \sin^2 \theta \sin^2 \varphi + \frac{^{(5)}\phi_3}{\xi} \left\{ 1 + \sin^2 \theta (2 \sin^2 \varphi - 5 \cos^2 \varphi) \right\} \right], \quad (4.32)$$

³For the $n = 0$ case, i.e., the incompressible fluid case, $^{(4)}\phi_2$ is proportional to ξ^2 .

$$\frac{\partial \tilde{\Phi}_6}{\partial \tilde{x}_1} = 30 \left[\left(\frac{d^{(6)}\phi_4}{d\xi} - \frac{4^{(6)}\phi_4}{\xi} \right) (3 - 7 \sin^2 \theta \cos^2 \varphi) \sin^3 \theta \sin \varphi \cos^2 \varphi + \frac{3^{(6)}\phi_4}{\xi} (1 - 5 \sin^2 \theta \cos^2 \varphi) \sin \theta \sin \varphi \right], \quad (4.33)$$

$$\begin{aligned} \frac{\partial \tilde{\Phi}_6}{\partial \tilde{x}_2} = 30 \left[\left(\frac{d^{(6)}\phi_4}{d\xi} - \frac{4^{(6)}\phi_4}{\xi} \right) (3 - 7 \sin^2 \theta \cos^2 \varphi) \sin^3 \theta \sin^2 \varphi \cos \varphi \right. \\ \left. + \frac{6^{(6)}\phi_4}{\xi} \left\{ 3 + \sin^2 \theta (7 \cos^2 \varphi - 6 \sin^2 \varphi) \right\} \sin \theta \cos \varphi \right]. \end{aligned} \quad (4.34)$$

In Table XIII, we show the first zero point of Θ_0 , the orbital separation at the contact point, the energies, the quadrupole moment, and the change in the central density. In this table, we have used

$$\tilde{E} \equiv \frac{E}{M^2/a_0} = \tilde{E}_{self} + \epsilon \tilde{E}_{point} + \epsilon^6 \tilde{E}_{quad}. \quad (4.35)$$

Each energy is given in Table XIV, where we have defined as

$$\tilde{\Pi}_{tot} = \tilde{\Pi}_{self} + \epsilon^6 \tilde{\Pi}_{quad}, \quad (4.36)$$

$$(\tilde{W}_{self})_{tot} = (\tilde{W}_{self})_{self} + \epsilon^6 (\tilde{W}_{self})_{quad}, \quad (4.37)$$

$$(\tilde{W}_{int})_{tot} = \epsilon (\tilde{W}_{int})_{point} + \epsilon^6 (\tilde{W}_{int})_{quad}, \quad (4.38)$$

$$\tilde{T}_{tot} = \epsilon \tilde{T}_{point} + \epsilon^6 \tilde{T}_{quad}. \quad (4.39)$$

Accordingly, we can express

$$\tilde{E}_{self} = \tilde{\Pi}_{self} + (\tilde{W}_{self})_{self}, \quad (4.40)$$

$$\tilde{E}_{point} = (\tilde{W}_{int})_{point} + \tilde{T}_{point}, \quad (4.41)$$

$$\tilde{E}_{quad} = \tilde{\Pi}_{quad} + (\tilde{W}_{self})_{quad} + (\tilde{W}_{int})_{quad} + \tilde{T}_{quad}. \quad (4.42)$$

The total angular momentum for the identical star binary becomes

$$J = \frac{M}{2} R^2 \Omega, \quad (4.43)$$

and the orbital angular velocity is calculated as

$$\Omega^2 = \frac{2M}{a_0^3} \epsilon^3 \left[1 + \frac{9\epsilon^5}{a_0^2} \frac{\tilde{T}_{11}}{M} + O(\epsilon^7) \right]. \quad (4.44)$$

We show the total energy, the total angular momentum and the orbital angular velocity along the equilibrium sequences of the binary system in Table XV.

In Figs. 1–4, we show (a) the total energy and (b) the total angular momentum as functions of the orbital separation, and (c) the orbital angular velocity as functions of the total angular momentum for each polytropic index, where we have used

$$\rho_0 \equiv \frac{3M}{4\pi a_0^3} \quad (4.45)$$

In these figures, we compare our results (solid lines) with those of numerical calculations computed by Uryū and Eriguchi [25] (open triangles) and those of semi-analytic ones proposed by Lai, Rasio and Shapiro [26] (filled circles). For the smaller orbital separation, we must expand the physical values up to higher order than $O(\epsilon^6)$ in order to include the effect of the spin of each star. In the case of the irrotational binary system, there is no intrinsic spin for the large distant stars. However, when the stars reach close range, the small spins are produced by the deformations of the stars. For example, the effect of these spins appears in the total energy at $O(\epsilon^9)^4$. The effect of the spin in the case of the smaller polytropic index is larger than that in the case of the larger polytropic index. The reason is

⁴This is because the spin kinetic energy is given by the volume integral of $(\epsilon^4 \nabla \Phi_4)^2 \sim \Omega^2 \epsilon^6 (\tilde{\nabla} \tilde{\Phi}_4)^2$, where Ω^2 is $O(\epsilon^3)$. The definition of $\tilde{\nabla}$ is in Eq. (A4).

that the matter is not concentrated in the case of the smaller polytropic index, and the quadrupole moment becomes larger (see Table XIII). Therefore, at the smaller orbital separation, the deviation between our results and those of Uryū and Eriguchi for $n = 0.5$ (Fig. 1) is larger than that for $n = 1.5$ (Fig. 3).

Finally, we represent the accuracy of our numerical calculations by comparing numerical and analytic solutions for $n = 1$ cases [29]. In Fig. 5, we show the absolute values of the relative error for $^{(3)}\psi_2$, $^{(4)}\psi_3$, $^{(5)}\psi_4$, $^{(6)}\psi_2$, $^{(6)}\psi_4$ and $^{(6)}\psi_5$, where we define the relative error by

$$\left| \frac{\text{Numerical solution}}{\text{Analytic solution}} - 1 \right|. \quad (4.46)$$

We can see from Fig. 5 that the relative errors are within 10^{-8} . Therefore, we regard that our results have enough accuracy.

V. SUMMARY AND DISCUSSIONS

A. Summary

In this paper, we have calculated the equilibrium solutions of irrotational binary polytropic stars in Newtonian gravity by expanding all physical quantities in a power of ϵ . We have presented the results of the cases of several polytropic indices ($n = 0.5, 1, 1.5$, and 2). In particular, we have shown the velocity fields by solving the equation of continuity. It is found that there exists the small velocity component along the orbital axis (see §IV). It agrees with the numerical calculations performed by Uryū and Eriguchi [25] and Bonazzola, Gourgoulhon and Marck [18,19,33].

Furthermore, we have given the figures and tables of the total energy, total angular momentum and orbital angular velocity along the equilibrium sequences for each polytropic indices. We can see from these figures that our results agree with those of Lai, Rasio and Shapiro [26] and Uryū and Eriguchi [25] for $R/a_0 > 3$.

Since our solutions are correct if $\epsilon \ll 1$, they can be used to check the validity of numerical solutions. For any numerical codes, one can ask to solve an equilibrium for large R , and compare numerically derived velocity distribution and so on with our semi-analytic solutions.

However, since we expanded physical quantities up to $O(\epsilon^6)$, it may not be enough to discuss about the behavior of solutions for small R . In order to apply our solutions in the case of small R and to check the validity of numerical codes in this case, further higher order calculations are needed.

B. Discussions

It is important to compare the velocity field which we obtain by solving the continuity equation with that given by Lai, Rasio and Shapiro. We can see the velocity field they give in their papers [26,32] or in the Chandrasekhar's textbook [34]. In the irrotational case, it becomes

$$(u_{LRS})_1 = \frac{2a_1^2}{a_1^2 + a_2^2} \Omega x_2, \quad (5.1)$$

$$(u_{LRS})_2 = -\frac{2a_2^2}{a_1^2 + a_2^2} \Omega x_1, \quad (5.2)$$

$$(u_{LRS})_3 = 0, \quad (5.3)$$

in the corotating frame. Here a_i denotes the length of the principal axis parallel to the x_i -axis. We can rewrite these component of the velocity field as

$$(u_{LRS})_1 = \Omega x_2 + \left(\frac{a_1^2 - a_2^2}{a_1^2 + a_2^2} \right) \Omega x_2, \quad (5.4)$$

$$(u_{LRS})_2 = -\Omega x_1 + \left(\frac{a_1^2 - a_2^2}{a_1^2 + a_2^2} \right) \Omega x_1. \quad (5.5)$$

The second terms in the above equations are order ϵ^3 because the deviation between a_1 and a_2 is produced by the tidal force and the effect of the tidal force is order ϵ^3 .

On the other hand, if we restrict the form of the function in our velocity field as ${}^{(4)}\phi_2 = (b_2/\xi_1)\xi^2$, ${}^{(5)}\phi_3 = (\alpha b_3/\xi_1)\xi^3$ and ${}^{(6)}\phi_4 = (\alpha^2 b_4/\xi_1)\xi^4$ as in the case of the incompressible fluid, we can express our velocity field by using Eqs. (4.25), (4.26) and (4.18) as

$$(u_{\text{present}})_1 = \Omega x_2 + \Omega [6\epsilon^3 b_2 x_2 - 48\epsilon^4 b_3 x_1 x_2 + 90\epsilon^5 b_4 x_2 (r^2 - 5x_1^2)], \quad (5.6)$$

$$(u_{\text{present}})_2 = -\Omega x_1 + \Omega [6\epsilon^3 b_2 x_1 + 6\epsilon^4 b_3 (r^2 - 5x_1^2 + 2x_2^2) + 30\epsilon^5 b_4 r (3r^2 + 7x_1^2 - 6x_2^2)], \quad (5.7)$$

$$(u_{\text{present}})_3 = \Omega [12\epsilon^4 b_3 x_2 x_3 + 180\epsilon^5 b_4 x_1 x_2 x_3], \quad (5.8)$$

where $r^2 = x_1^2 + x_2^2 + x_3^2$. Therefore, we find that the form of the velocity field we obtain in the case of the incompressible fluid coincides with that given by Lai, Rasio and Shapiro up to $O(\epsilon^3)$ including the value of b_2 . This means that the velocity field of Lai, Rasio and Shapiro is correct only in the case of the “incompressible” equation of state and “ellipsoidal” figures, because the velocity field at order ϵ^3 is produced by the ellipsoidal deformation of star 1 (see Appendix F).

Finally, we discuss about the configuration of each star. When we pay attention to star 1, the equation for the stellar surface is written as

$$\begin{aligned} \Xi(\theta, \varphi) &= \xi_1 + \epsilon^3 S_3(\theta, \varphi) + \epsilon^4 S_4(\theta, \varphi) + \epsilon^5 S_5(\theta, \varphi) + \epsilon^6 S_6(\theta, \varphi), \\ &= \xi_1 + \epsilon^3 \frac{{}^{(3)}\psi_2(\xi_1)}{|\Theta_{0,\xi}(\xi_1)|} P_2(\sin \theta \cos \varphi) + \epsilon^4 \frac{{}^{(4)}\psi_3(\xi_1)}{|\Theta_{0,\xi}(\xi_1)|} P_3(\sin \theta \cos \varphi) + \epsilon^5 \frac{{}^{(5)}\psi_4(\xi_1)}{|\Theta_{0,\xi}(\xi_1)|} P_4(\sin \theta \cos \varphi) \\ &\quad + \epsilon^6 \frac{1}{|\Theta_{0,\xi}(\xi_1)|} \left[\frac{{}^{(3)}\psi_2(\xi_1)}{|\Theta_{0,\xi}(\xi_1)|} \left(\frac{{}^{(3)}\psi_2(\xi_1)}{\xi_1} + \frac{d({}^{(3)}\psi_2)}{d\xi}(\xi_1) \right) \left\{ \frac{18}{35} P_4(\sin \theta \cos \varphi) + \frac{2}{7} P_2(\sin \theta \cos \varphi) + \frac{1}{5} \right\} \right. \\ &\quad \left. + {}^{(6)}\psi_0(\xi_1) + {}^{(6)}\psi_2(\xi_1) P_2(\sin \theta \cos \varphi) + {}^{(6)}\psi_{22}(\xi_1) P_2^2(\cos \theta) \cos 2\varphi \right. \\ &\quad \left. + {}^{(6)}\psi_4(\xi_1) P_4(\sin \theta \cos \varphi) + {}^{(6)}\psi_5(\xi_1) P_5(\sin \theta \cos \varphi) \right], \end{aligned} \quad (5.10)$$

where S_i are defined in Appendix A and we have used the relation (C27). Accordingly, we can express the length of the principal axis. Although the real length of the axis is written as $\alpha\Xi$, we show the results divided by α .

(1) The opposite direction to star 2:

$$\begin{aligned} \Xi\left(\frac{\pi}{2}, 0\right) &= \xi_1 + \epsilon^3 \frac{{}^{(3)}\psi_2(\xi_1)}{|\Theta_{0,\xi}(\xi_1)|} + \epsilon^4 \frac{{}^{(4)}\psi_3(\xi_1)}{|\Theta_{0,\xi}(\xi_1)|} + \epsilon^5 \frac{{}^{(5)}\psi_4(\xi_1)}{|\Theta_{0,\xi}(\xi_1)|} \\ &\quad + \epsilon^6 \frac{1}{|\Theta_{0,\xi}(\xi_1)|} \left[\frac{{}^{(3)}\psi_2(\xi_1)}{|\Theta_{0,\xi}(\xi_1)|} \left(\frac{{}^{(3)}\psi_2(\xi_1)}{\xi_1} + \frac{d({}^{(3)}\psi_2)}{d\xi}(\xi_1) \right) + {}^{(6)}\psi_0(\xi_1) + {}^{(6)}\psi_2(\xi_1) + 3{}^{(6)}\psi_{22}(\xi_1) \right. \\ &\quad \left. + {}^{(6)}\psi_4(\xi_1) + {}^{(6)}\psi_5(\xi_1) \right], \end{aligned} \quad (5.11)$$

(2) The direction to star 2:

$$\begin{aligned} \Xi\left(\frac{\pi}{2}, \pi\right) &= \xi_1 + \epsilon^3 \frac{{}^{(3)}\psi_2(\xi_1)}{|\Theta_{0,\xi}(\xi_1)|} - \epsilon^4 \frac{{}^{(4)}\psi_3(\xi_1)}{|\Theta_{0,\xi}(\xi_1)|} + \epsilon^5 \frac{{}^{(5)}\psi_4(\xi_1)}{|\Theta_{0,\xi}(\xi_1)|} \\ &\quad + \epsilon^6 \frac{1}{|\Theta_{0,\xi}(\xi_1)|} \left[\frac{{}^{(3)}\psi_2(\xi_1)}{|\Theta_{0,\xi}(\xi_1)|} \left(\frac{{}^{(3)}\psi_2(\xi_1)}{\xi_1} + \frac{d({}^{(3)}\psi_2)}{d\xi}(\xi_1) \right) + {}^{(6)}\psi_0(\xi_1) + {}^{(6)}\psi_2(\xi_1) + 3{}^{(6)}\psi_{22}(\xi_1) \right. \\ &\quad \left. + {}^{(6)}\psi_4(\xi_1) - {}^{(6)}\psi_5(\xi_1) \right], \end{aligned} \quad (5.12)$$

(3) The (opposite) direction to the orbital motion:

$$\begin{aligned} \Xi\left(\frac{\pi}{2}, \frac{\pi}{2}\right) &= \Xi\left(\frac{\pi}{2}, \frac{3\pi}{2}\right), \\ &= \xi_1 - \epsilon^3 \frac{{}^{(3)}\psi_2(\xi_1)}{2|\Theta_{0,\xi}(\xi_1)|} + \epsilon^5 \frac{3{}^{(5)}\psi_4(\xi_1)}{8|\Theta_{0,\xi}(\xi_1)|} \\ &\quad + \epsilon^6 \frac{1}{|\Theta_{0,\xi}(\xi_1)|} \left[\frac{{}^{(3)}\psi_2(\xi_1)}{4|\Theta_{0,\xi}(\xi_1)|} \left(\frac{{}^{(3)}\psi_2(\xi_1)}{\xi_1} + \frac{d({}^{(3)}\psi_2)}{d\xi}(\xi_1) \right) + {}^{(6)}\psi_0(\xi_1) - \frac{1}{2} {}^{(6)}\psi_2(\xi_1) - 3{}^{(6)}\psi_{22}(\xi_1) \right. \\ &\quad \left. + \frac{3}{8} {}^{(6)}\psi_4(\xi_1) \right], \end{aligned} \quad (5.13)$$

(4) The direction parallel to the rotational axis:

$$\begin{aligned}\Xi(0, 0) &= \Xi(-\pi, 0), \\ &= \xi_1 - \epsilon^3 \frac{{}^{(3)}\psi_2(\xi_1)}{2|\Theta_{0,\xi}(\xi_1)|} + \epsilon^5 \frac{3{}^{(5)}\psi_4(\xi_1)}{8|\Theta_{0,\xi}(\xi_1)|} \\ &\quad + \epsilon^6 \frac{1}{|\Theta_{0,\xi}(\xi_1)|} \left[\frac{{}^{(3)}\psi_2(\xi_1)}{4|\Theta_{0,\xi}(\xi_1)|} \left(\frac{{}^{(3)}\psi_2(\xi_1)}{\xi_1} + \frac{d{}^{(3)}\psi_2}{d\xi}(\xi_1) \right) + {}^{(6)}\psi_0(\xi_1) - \frac{1}{2}{}^{(6)}\psi_2(\xi_1) + \frac{3}{8}{}^{(6)}\psi_4(\xi_1) \right].\end{aligned}\quad (5.14)$$

We can see from these equations and Tables I – VIII that the axis to star 2 is the longest, and the deviation between the axis to star 2 and that opposite to star 2 appears at order ϵ^4 . On the contrary, the deviation between the axis to the orbital motion and that parallel to the rotational axis appears at order ϵ^6 , and the difference is the effect of the deformation induced by the spin of the figure (${}^{(6)}\psi_{22}$).

When we see the quadrupole moments in Eq. (5.10), we find that the coefficient of the higher order term is not large as

$$\epsilon^3 \frac{{}^{(3)}\psi_2(\xi_1)}{|\Theta_{0,\xi}(\xi_1)|} \left[1 + \epsilon^3 \left\{ \frac{2}{7|\Theta_{0,\xi}(\xi_1)|} \left(\frac{{}^{(3)}\psi_2(\xi_1)}{\xi_1} + \frac{d{}^{(3)}\psi_2}{d\xi}(\xi_1) \right) + \frac{{}^{(6)}\psi_2(\xi_1)}{{}^{(3)}\psi_2(\xi_1)} \right\} \right] P_2(\sin \theta \cos \varphi). \quad (5.15)$$

Therefore we can expect the convergence of these terms. However, there is another quadrupole moment induced by the spin of the figure. The term $\epsilon^6 ({}^{(6)}\psi_{22}(\xi_1)/|\Theta_{0,\xi}(\xi_1)|) P_2^2(\cos \theta) \cos 2\varphi$ seems to be as effective as the leading quadrupole term for the smaller orbital separation. This means that the terms concerned with the spin of the figure which appear at order ϵ^9 in the total energy may change the behavior of the total energy. Furthermore, the hexadecapole moments in Eq. (5.10),

$$\epsilon^5 \frac{{}^{(5)}\psi_4(\xi_1)}{|\Theta_{0,\xi}(\xi_1)|} \left[1 + \epsilon \left\{ \frac{18}{35{}^{(5)}\psi_4(\xi_1)} \frac{{}^{(3)}\psi_2(\xi_1)}{|\Theta_{0,\xi}(\xi_1)|} \left(\frac{{}^{(3)}\psi_2(\xi_1)}{\xi_1} + \frac{d{}^{(3)}\psi_2}{d\xi}(\xi_1) \right) + \frac{{}^{(6)}\psi_4(\xi_1)}{{}^{(5)}\psi_4(\xi_1)} \right\} \right] P_4(\sin \theta \cos \varphi), \quad (5.16)$$

does not seem to converge at order ϵ^6 . However, since these terms will appear at order ϵ^{10} in the total energy, they do not have so much effect. Anyway, if we discuss about the behavior of the total energy, the total angular momentum and so on for the smaller orbital separation such as $R/a_0 < 3$, we must calculate at least up to order ϵ^9 .

ACKNOWLEDGMENTS

We would like to thank K. Ioka and K. Nakao for useful discussions. This work was partly supported by a Grant-in-Aid for Scientific Research Fellowship (No.9402: KT) and Grant-in-Aid of Scientific Research (No.11640274, 09NP0801: TN) of the Japanese Ministry of Education, Science, Sports and Culture.

APPENDIX A: SUMMARY OF THE EQUATIONS AND THEIR BOUNDARY CONDITIONS

In the following appendices, we derive equations order by order of ϵ , and describe in detail. In this appendix, we summarize the equations which should be solved numerically with their boundary conditions in each order.

First of all, we give the total equations which include all terms up to $O(\epsilon^6)$. The equations for determination of the velocity potential and stellar configuration are written as

$$\tilde{\Delta}\tilde{\Phi} = -n \left[\tilde{\nabla}\tilde{\Phi} - (\tilde{\Omega} \times \boldsymbol{\xi})_{orb} - \frac{\epsilon}{\xi_1} (\tilde{\Omega} \times \boldsymbol{\xi})_{fig} \right] \cdot \frac{\tilde{\nabla}\Theta}{\Theta}, \quad (A1)$$

$$\tilde{\Delta}\Theta = -\Theta^n - \frac{\Omega^2 \xi_1^2}{8\pi\rho_c \epsilon^2} \tilde{\Delta} \left[(\tilde{\nabla}\tilde{\Phi})^2 - 2(\tilde{\nabla}\tilde{\Phi}) \cdot \left\{ (\tilde{\Omega} \times \boldsymbol{\xi})_{orb} + \frac{\epsilon}{\xi_1} (\tilde{\Omega} \times \boldsymbol{\xi})_{fig} \right\} \right], \quad (A2)$$

where

$$\tilde{\Delta} = \alpha^2 \Delta = \frac{1}{\xi^2} \frac{\partial}{\partial \xi} \left(\xi^2 \frac{\partial}{\partial \xi} \right) + \frac{1}{\xi^2 \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial}{\partial \theta} \right) + \frac{1}{\xi^2 \sin^2 \theta} \frac{\partial^2}{\partial \varphi^2}, \quad (A3)$$

$$\tilde{\nabla} = \alpha \nabla = \frac{\partial}{\partial \xi} \hat{\boldsymbol{\xi}} + \frac{1}{\xi} \frac{\partial}{\partial \theta} \hat{\boldsymbol{\theta}} + \frac{1}{\xi \sin \theta} \frac{\partial}{\partial \varphi} \hat{\boldsymbol{\varphi}}. \quad (A4)$$

The boundary condition for the velocity field is

$$\left[\tilde{\nabla} \tilde{\Phi} - (\tilde{\Omega} \times \boldsymbol{\xi})_{orb} - \frac{\epsilon}{\xi_1} (\tilde{\Omega} \times \boldsymbol{\xi})_{fig} \right] \cdot (\tilde{\nabla} \Theta) \Big|_{surf} = 0. \quad (\text{A5})$$

The internal and external gravitational potentials which should be matched at the stellar surface are

$$\begin{aligned} \tilde{U}_{int} &\equiv \frac{U^{1 \rightarrow 1}}{4\pi\rho_c\alpha^2} \\ &= \Theta + \frac{\Omega^2 \xi_1^2}{8\pi\rho_c\epsilon^2} \left[(\tilde{\nabla} \tilde{\Phi})^2 - 2(\tilde{\nabla} \tilde{\Phi}) \cdot \left\{ (\tilde{\Omega} \times \boldsymbol{\xi})_{orb} + \frac{\epsilon}{\xi_1} (\tilde{\Omega} \times \boldsymbol{\xi})_{fig} \right\} \right] + \tilde{U}_0 \\ &\quad - \frac{\mu}{1+p} \epsilon \sum_{l=0}^{\infty} (-1)^l \epsilon^l \left(\frac{\xi}{\xi_1} \right)^l P_l(\sin \theta \cos \varphi) - \frac{3\mu}{2M_{tot}} \left(\frac{I'_{11}}{a_0^2} \right) \epsilon^3 \left[1 - 3\epsilon \frac{\xi}{\xi_1} P_1(\sin \theta \cos \varphi) + \dots \right], \end{aligned} \quad (\text{A6})$$

$$\begin{aligned} \tilde{U}_{ext} &= \frac{\kappa_0}{\xi} + \epsilon \sum_{l,m} \frac{\kappa_{1:lm}}{\xi^{l+1}} Y_l^m + \epsilon^2 \sum_{l,m} \frac{\kappa_{2:lm}}{\xi^{l+1}} Y_l^m + \epsilon^3 \sum_{l,m} \frac{\kappa_{3:lm}}{\xi^{l+1}} Y_l^m + \epsilon^4 \sum_{l,m} \frac{\kappa_{4:lm}}{\xi^{l+1}} Y_l^m + \epsilon^5 \sum_{l,m} \frac{\kappa_{5:lm}}{\xi^{l+1}} Y_l^m \\ &\quad + \epsilon^6 \sum_{l,m} \frac{\kappa_{6:lm}}{\xi^{l+1}} Y_l^m + \dots, \end{aligned} \quad (\text{A7})$$

where κ_0 and $\kappa_{i:lm}$ are multipole moments. $\tilde{U}_0$ is constant and expanded as

$$\tilde{U}_0 = c_0 + \epsilon c_1 + \epsilon^2 c_2 + \epsilon^3 c_3 + \epsilon^4 c_4 + \epsilon^5 c_5 + \epsilon^6 c_6. \quad (\text{A8})$$

The velocity potential and configuration function are expanded up to $O(\epsilon^6)$ as

$$\tilde{\Phi} = \tilde{\Phi}_0 + \epsilon \tilde{\Phi}_1 + \epsilon^2 \tilde{\Phi}_2 + \epsilon^3 \tilde{\Phi}_3 + \epsilon^4 \tilde{\Phi}_4 + \epsilon^5 \tilde{\Phi}_5 + \epsilon^6 \tilde{\Phi}_6, \quad (\text{A9})$$

$$\Theta = \Theta_0 + \epsilon \Theta_1 + \epsilon^2 \Theta_2 + \epsilon^3 \Theta_3 + \epsilon^4 \Theta_4 + \epsilon^5 \Theta_5 + \epsilon^6 \Theta_6. \quad (\text{A10})$$

The boundary conditions for Θ are

$$\tilde{U}_{int}(\Xi) = \tilde{U}_{ext}(\Xi), \quad (\text{A11})$$

$$\frac{\partial \tilde{U}_{int}}{\partial \xi}(\Xi) = \frac{\partial \tilde{U}_{ext}}{\partial \xi}(\Xi), \quad (\text{A12})$$

where Ξ denotes the first zero point of the function Θ , i.e., the stellar surface and is formally expressed as

$$\Xi(\theta, \varphi) = \xi_1 + \epsilon S_1(\theta, \varphi) + \epsilon^2 S_2(\theta, \varphi) + \epsilon^3 S_3(\theta, \varphi) + \epsilon^4 S_4(\theta, \varphi) + \epsilon^5 S_5(\theta, \varphi) + \epsilon^6 S_6(\theta, \varphi). \quad (\text{A13})$$

The regularity conditions at the center of each star is

$$\frac{\partial \Theta}{\partial \xi}(\xi = 0) = 0, \quad (\text{A14})$$

and also we normalize the central value of Θ as

$$\Theta(\xi = 0) = 1. \quad (\text{A15})$$

In the following, we show the equation for the velocity potential, its boundary condition, the equation for determination of the figure and its boundary conditions (A11) and (A12).

1. 0th order

The equation for the velocity potential is

$$\tilde{\Delta} \tilde{\Phi}_0 = -n \left[\tilde{\nabla} \tilde{\Phi}_0 - (\tilde{\Omega} \times \boldsymbol{\xi})_{orb} \right] \cdot \frac{\tilde{\nabla} \Theta_0}{\Theta_0}, \quad (\text{A16})$$

and its boundary condition is

$$\left[\tilde{\nabla} \tilde{\Phi}_0 - (\tilde{\Omega} \times \boldsymbol{\xi})_{orb} \right] \cdot (\tilde{\nabla} \Theta_0) \Big|_{\xi_1} = 0. \quad (\text{A17})$$

The equation for determination of the figure is

$$\frac{1}{\xi^2} \frac{d}{d\xi} \left(\xi^2 \frac{d\Theta_0}{d\xi} \right) = -\Theta_0^n, \quad (\text{A18})$$

and its boundary conditions at the stellar surface are

$$c_0 = \frac{\kappa_0}{\xi_1}, \quad (\text{A19})$$

$$\frac{d\Theta_0}{d\xi}(\xi_1) = -\frac{\kappa_0}{\xi_1^2}. \quad (\text{A20})$$

The regularity condition at the center of the star is

$$\frac{d\Theta_0}{d\xi}(\xi = 0) = 0, \quad (\text{A21})$$

and the normalization of Θ_0 at the center of the star is

$$\Theta_0(\xi = 0) = 1. \quad (\text{A22})$$

2. 1st order

The equation for the velocity potential is

$$\tilde{\Delta} \tilde{\Phi}_1 = -n \left[\tilde{\nabla} \tilde{\Phi}_1 - \frac{1}{\xi_1} (\tilde{\Omega} \times \boldsymbol{\xi})_{fig} \right] \cdot \frac{\tilde{\nabla} \Theta_0}{\Theta_0}, \quad (\text{A23})$$

and its boundary condition is

$$(\tilde{\nabla} \tilde{\Phi}_1) \cdot (\tilde{\nabla} \Theta_0) \Big|_{\xi_1} = 0. \quad (\text{A24})$$

The equation for determination of the figure is

$$\tilde{\Delta} \Theta_1 = -n \Theta_0^{n-1} \Theta_1, \quad (\text{A25})$$

and its boundary conditions at the stellar surface are

$$c_1 - \frac{\mu(3+2p)}{2(1+p)^2} = -\frac{\kappa_0}{\xi_1^2} S_1 + \sum_{l,m} \frac{\kappa_{1:lm}}{\xi_1^{l+1}} Y_l^m, \quad (\text{A26})$$

$$S_1 \frac{d^2 \Theta_0}{d\xi^2}(\xi_1) + \frac{\partial \Theta_1}{\partial \xi}(\xi_1) = \frac{2\kappa_0}{\xi_1^3} S_1 - \sum_{l,m} (l+1) \frac{\kappa_{1:lm}}{\xi_1^{l+2}} Y_l^m. \quad (\text{A27})$$

The regularity condition at the center of the star is

$$\frac{\partial \Theta_1}{\partial \xi}(\xi = 0) = 0, \quad (\text{A28})$$

and the normalization of Θ_1 at the center of the star is

$$\Theta_1(\xi = 0) = 0, \quad (\text{A29})$$

because we take $\Theta_0(\xi = 0) = 1$.

3. 2nd order

The equation for the velocity potential is

$$\tilde{\Delta}\tilde{\Phi}_2 = -n(\tilde{\nabla}\tilde{\Phi}_2) \cdot \frac{\tilde{\nabla}\Theta_0}{\Theta_0}, \quad (\text{A30})$$

and its boundary condition is

$$(\tilde{\nabla}\tilde{\Phi}_2) \cdot (\tilde{\nabla}\Theta_0)|_{\xi_1} = 0. \quad (\text{A31})$$

The equation for determination of the figure is

$$\tilde{\Delta}\Theta_2 = -n\Theta_0^{n-1}\Theta_2, \quad (\text{A32})$$

and its boundary conditions at the stellar surface are

$$c_2 = -\frac{\kappa_0}{\xi_1^2}S_2 + \sum_{l,m} \frac{\kappa_{2:lm}}{\xi_1^{l+1}}Y_l^m, \quad (\text{A33})$$

$$S_2 \frac{d^2\Theta_0}{d\xi^2}(\xi_1) + \frac{\partial\Theta_2}{\partial\xi}(\xi_1) = \frac{2\kappa_0}{\xi_1^3}S_2 - \sum_{l,m} (l+1) \frac{\kappa_{2:lm}}{\xi_1^{l+2}}Y_l^m. \quad (\text{A34})$$

The regularity condition at the center of the star is

$$\frac{\partial\Theta_2}{\partial\xi}(\xi=0) = 0, \quad (\text{A35})$$

and the normalization of Θ_2 at the center of the star is

$$\Theta_2(\xi=0) = 0, \quad (\text{A36})$$

because we take $\Theta_0(\xi=0) = 1$.

4. 3rd order

The equation for the velocity potential is

$$\tilde{\Delta}\tilde{\Phi}_3 = -n(\tilde{\nabla}\tilde{\Phi}_3) \cdot \frac{\tilde{\nabla}\Theta_0}{\Theta_0}, \quad (\text{A37})$$

and its boundary condition is

$$(\tilde{\nabla}\tilde{\Phi}_3) \cdot (\tilde{\nabla}\Theta_0)|_{\xi_1} = 0. \quad (\text{A38})$$

The equation for determination of the figure is

$$\tilde{\Delta}\Theta_3 = -n\Theta_0^{n-1}\Theta_3, \quad (\text{A39})$$

and its boundary conditions at the stellar surface are

$$c_3 - \frac{\mu}{1+p}P_2(\sin\theta\cos\varphi) = -\frac{\kappa_0}{\xi_1^2}S_3 + \sum_{l,m} \frac{\kappa_{3:lm}}{\xi_1^{l+1}}Y_l^m, \quad (\text{A40})$$

$$S_3 \frac{d^2\Theta_0}{d\xi^2}(\xi_1) + \frac{\partial\Theta_3}{\partial\xi}(\xi_1) - \frac{2\mu}{\xi_1(1+p)}P_2(\sin\theta\cos\varphi) = \frac{2\kappa_0}{\xi_1^3}S_3 - \sum_{l,m} (l+1) \frac{\kappa_{3:lm}}{\xi_1^{l+2}}Y_l^m. \quad (\text{A41})$$

The regularity condition at the center of the star is

$$\frac{\partial\Theta_3}{\partial\xi}(\xi=0) = 0, \quad (\text{A42})$$

and the normalization of Θ_3 at the center of the star is

$$\Theta_3(\xi=0) = 0, \quad (\text{A43})$$

because we take $\Theta_0(\xi=0) = 1$.

5. 4th order

The equation for the velocity potential is

$$\tilde{\Delta}\tilde{\Phi}_4 = -\frac{n}{\Theta_0} \left[(\tilde{\nabla}\tilde{\Phi}_4) \cdot (\tilde{\nabla}\Theta_0) - \frac{1}{\xi_1} (\tilde{\Omega} \times \boldsymbol{\xi})_{fig} \cdot (\tilde{\nabla}\Theta_3) \right], \quad (\text{A44})$$

and its boundary condition is

$$\left[(\tilde{\nabla}\tilde{\Phi}_4) \cdot (\tilde{\nabla}\Theta_0) - \frac{1}{\xi_1} (\tilde{\Omega} \times \boldsymbol{\xi})_{fig} \cdot (\tilde{\nabla}\Theta_3) \right] \Big|_{\xi_1} = 0. \quad (\text{A45})$$

The equation for determination of the figure is

$$\tilde{\Delta}\Theta_4 = -n\Theta_0^{n-1}\Theta_4, \quad (\text{A46})$$

and its boundary conditions at the stellar surface are

$$c_4 + \frac{\mu}{1+p} P_3(\sin\theta \cos\varphi) = -\frac{\kappa_0}{\xi_1^2} S_4 + \sum_{l,m} \frac{\kappa_{4:lm}}{\xi_1^{l+1}} Y_l^m, \quad (\text{A47})$$

$$S_4 \frac{d^2\Theta_0}{d\xi^2}(\xi_1) + \frac{\partial\Theta_4}{\partial\xi}(\xi_1) + \frac{3\mu}{\xi_1(1+p)} P_3(\sin\theta \cos\varphi) = \frac{2\kappa_0}{\xi_1^3} S_4 - \sum_{l,m} (l+1) \frac{\kappa_{4:lm}}{\xi_1^{l+2}} Y_l^m. \quad (\text{A48})$$

The regularity condition at the center of the star is

$$\frac{\partial\Theta_4}{\partial\xi}(\xi=0) = 0, \quad (\text{A49})$$

and the normalization of Θ_4 at the center of the star is

$$\Theta_4(\xi=0) = 0, \quad (\text{A50})$$

because we take $\Theta_0(\xi=0) = 1$.

6. 5th order

The equation for the velocity potential is

$$\tilde{\Delta}\tilde{\Phi}_5 = -\frac{n}{\Theta_0} \left[(\tilde{\nabla}\tilde{\Phi}_5) \cdot (\tilde{\nabla}\Theta_0) - \frac{1}{\xi_1} (\tilde{\Omega} \times \boldsymbol{\xi})_{fig} \cdot (\tilde{\nabla}\Theta_4) \right], \quad (\text{A51})$$

and its boundary condition is

$$\left[(\tilde{\nabla}\tilde{\Phi}_5) \cdot (\tilde{\nabla}\Theta_0) - \frac{1}{\xi_1} (\tilde{\Omega} \times \boldsymbol{\xi})_{fig} \cdot (\tilde{\nabla}\Theta_4) \right] \Big|_{\xi_1} = 0. \quad (\text{A52})$$

The equation for determination of the figure is

$$\tilde{\Delta}\Theta_5 = -n\Theta_0^{n-1}\Theta_5, \quad (\text{A53})$$

and its boundary conditions at the stellar surface are

$$c_5 - \frac{\mu}{1+p} P_4(\sin\theta \cos\varphi) = -\frac{\kappa_0}{\xi_1^2} S_5 + \sum_{l,m} \frac{\kappa_{5:lm}}{\xi_1^{l+1}} Y_l^m, \quad (\text{A54})$$

$$S_5 \frac{d^2\Theta_0}{d\xi^2}(\xi_1) + \frac{\partial\Theta_5}{\partial\xi}(\xi_1) - \frac{4\mu}{\xi_1(1+p)} P_4(\sin\theta \cos\varphi) = \frac{2\kappa_0}{\xi_1^3} S_5 - \sum_{l,m} (l+1) \frac{\kappa_{5:lm}}{\xi_1^{l+2}} Y_l^m. \quad (\text{A55})$$

The regularity condition at the center of the star is

$$\frac{\partial\Theta_5}{\partial\xi}(\xi=0) = 0, \quad (\text{A56})$$

and the normalization of Θ_5 at the center of the star is

$$\Theta_5(\xi=0) = 0, \quad (\text{A57})$$

because we take $\Theta_0(\xi=0) = 1$.

7. 6th order

The equation for the velocity potential is

$$\tilde{\Delta}\tilde{\Phi}_6 = -\frac{n}{\Theta_0} \left[(\tilde{\nabla}\tilde{\Phi}_6) \cdot (\tilde{\nabla}\Theta_0) - \frac{1}{\xi_1} (\tilde{\Omega} \times \boldsymbol{\xi})_{fig} \cdot (\tilde{\nabla}\Theta_5) \right], \quad (\text{A58})$$

and its boundary condition is

$$\left[(\tilde{\nabla}\tilde{\Phi}_6) \cdot (\tilde{\nabla}\Theta_0) - \frac{1}{\xi_1} (\tilde{\Omega} \times \boldsymbol{\xi})_{fig} \cdot (\tilde{\nabla}\Theta_5) \right] \Big|_{\xi_1} = 0. \quad (\text{A59})$$

The equation for determination of the figure is

$$\tilde{\Delta}\Theta_6 = -n\Theta_0^{n-1}\Theta_6 - \frac{1}{2}n(n-1)\Theta_0^{n-2}\Theta_3^2 + \frac{\mu}{\xi_1}\tilde{\Delta}\frac{\partial\tilde{\Phi}_4}{\partial\varphi}, \quad (\text{A60})$$

and its boundary conditions at the stellar surface are

$$\begin{aligned} c_6 - \frac{3\mu\bar{I}'_{11}}{2M_{tot}a_0^2} - \frac{\mu\delta}{2(1+p)^2} - \frac{2\mu}{\xi_1(1+p)}S_3P_2(\sin\theta\cos\varphi) + \frac{\mu}{1+p}P_5(\sin\theta\cos\varphi) - \frac{\mu}{\xi_1}\frac{\partial\tilde{\Phi}_4}{\partial\varphi}(\xi_1) \\ = \frac{\kappa_0}{\xi_1^2}\left(\frac{S_3^2}{\xi_1} - S_6\right) - \sum_{l,m}(l+1)\frac{\kappa_{3:lm}}{\xi_1^{l+2}}S_3Y_l^m + \sum_{l,m}\frac{\kappa_{6:lm}}{\xi_1^{l+1}}Y_l^m, \end{aligned} \quad (\text{A61})$$

$$\begin{aligned} S_6\frac{d^2\Theta_0}{d\xi^2}(\xi_1) + \frac{1}{2}S_3^2\frac{d^3\Theta_0}{d\xi^3}(\xi_1) + S_3\frac{\partial^2\Theta_3}{\partial\xi^2}(\xi_1) + \frac{\partial\Theta_6}{\partial\xi}(\xi_1) - \frac{2\mu}{\xi_1^2(1+p)}S_3P_2(\sin\theta\cos\varphi) + \frac{5\mu}{\xi_1(1+p)}P_5(\sin\theta\cos\varphi) \\ - \frac{\mu}{\xi_1}\frac{\partial^2\tilde{\Phi}_4}{\partial\xi\partial\varphi}(\xi_1) = -\frac{\kappa_0}{\xi_1^3}\left(\frac{3S_3^2}{\xi_1} - 2S_6\right) + \sum_{l=0}^{\infty}(l+1)(l+2)\frac{\kappa_{3:lm}}{\xi_1^{l+3}}S_3Y_l^m - \sum_{l=0}^{\infty}(l+1)\frac{\kappa_{6:lm}}{\xi_1^{l+2}}Y_l^m, \end{aligned} \quad (\text{A62})$$

where

$$\delta \equiv \frac{9}{2a_0^2} \left[\frac{\bar{I}_{11}}{M_1} + \left(\frac{a'_0}{a_0} \right)^3 \frac{\bar{I}'_{11}}{M_2} \right]. \quad (\text{A63})$$

The regularity condition at the center of the star is

$$\frac{\partial\Theta_6}{\partial\xi}(\xi=0) = 0, \quad (\text{A64})$$

and the normalization of Θ_6 at the center of the star is

$$\Theta_6(\xi=0) = 0, \quad (\text{A65})$$

because we take $\Theta_0(\xi=0) = 1$.

APPENDIX B: SOLUTIONS AT ZEROth ORDER

Since the configuration of star 1 is spherical at 0th order, the equation for determination of the figure becomes *Lane-Emden equation*,

$$\frac{1}{\xi^2} \frac{d}{d\xi} \left(\xi^2 \frac{d\Theta_0}{d\xi} \right) = -\Theta_0^n. \quad (\text{B1})$$

For example, we obtain analytic solutions,

$$\Theta_0 = 1 - \frac{\xi^2}{6} \quad (\text{for } n=0), \quad (\text{B2})$$

$$\Theta_0 = \frac{\sin\xi}{\xi} \quad (\text{for } n=1), \quad (\text{B3})$$

or numerical solutions by solving the ordinary differential equation (B1).

The velocity field at 0th order is the same as the orbital motion so that the velocity potential is determined by

$$\nabla\Phi_0 = \Omega R(\tilde{\Omega} \times \xi)_{orb}. \quad (B4)$$

Then, the solution should be

$$\Phi_0 = \frac{\Omega R}{1+p} r \sin \theta \sin \varphi. \quad (B5)$$

From this equation, we find that it is convenient to normalize Φ_0 as

$$\tilde{\Phi}_0 = \frac{1}{1+p} \xi \sin \theta \sin \varphi. \quad (B6)$$

In the following, we use the normalized velocity potential defined by

$$\tilde{\Phi} = \frac{\epsilon \Phi}{\Omega \alpha a_0}. \quad (B7)$$

Here we show the useful relations:

$$(\tilde{\nabla} \tilde{\Phi}_0)^2 = \frac{1}{(1+p)^2}, \quad (B8)$$

$$(\tilde{\nabla} \tilde{\Phi}_0) \cdot (\tilde{\Omega} \times \xi)_{orb} = \frac{1}{(1+p)^2}, \quad (B9)$$

$$(\tilde{\nabla} \tilde{\Phi}_0) \cdot (\tilde{\Omega} \times \xi)_{fig} = \frac{1}{1+p} \xi \sin \theta \cos \varphi. \quad (B10)$$

The orbital angular velocity can be calculated from Eq. (2.28) as

$$0 = \int_{star} d^3x \rho \left(-\frac{M_2}{R^2} + \frac{\Omega_0^2 R}{1+p} \right). \quad (B11)$$

Therefore, we have

$$\Omega_0^2 = \frac{M_{tot}}{R^3}, \quad (B12)$$

where $M_{tot} = M_1 + M_2$.

For 0th order, we have the internal and external gravitational potentials and their derivatives as

$$\tilde{U}_{int}(\xi) = \Theta_0 + c_0, \quad (B13)$$

$$\frac{\partial \tilde{U}_{int}}{\partial \xi}(\xi) = \frac{d\Theta_0}{d\xi}, \quad (B14)$$

$$\tilde{U}_{ext}(\xi) = \frac{\kappa_0}{\xi}, \quad (B15)$$

$$\frac{\partial \tilde{U}_{ext}}{\partial \xi}(\xi) = -\frac{\kappa_0}{\xi^2}. \quad (B16)$$

They become at the stellar surface as

$$\tilde{U}_{int}(\xi_1) = c_0, \quad (B17)$$

$$\frac{\partial \tilde{U}_{int}}{\partial \xi}(\xi_1) = \frac{d\Theta_0}{d\xi}(\xi_1), \quad (B18)$$

$$\tilde{U}_{ext}(\xi_1) = \frac{\kappa_0}{\xi_1}, \quad (B19)$$

$$\frac{\partial \tilde{U}_{ext}}{\partial \xi}(\xi_1) = -\frac{\kappa_0}{\xi_1^2}. \quad (B20)$$

Therefore, we can decide two constants at 0th order from the boundary conditions at the stellar surface as

$$\kappa_0 = -\xi_1^2 \frac{d\Theta_0}{d\xi}(\xi_1), \quad (\text{B21})$$

$$c_0 = -\xi_1 \frac{d\Theta_0}{d\xi}(\xi_1). \quad (\text{B22})$$

APPENDIX C: SOLUTIONS AT FIRST ORDER

The equation for the velocity potential is

$$\tilde{\Delta}\tilde{\Phi}_1 = -n \left[(\tilde{\nabla}\tilde{\Phi}_1) - \frac{1}{\xi_1} (\tilde{\Omega} \times \boldsymbol{\xi})_{fig} \right] \cdot \frac{\tilde{\nabla}\Theta_0}{\Theta_0}, \quad (\text{C1})$$

$$= -n (\tilde{\nabla}\tilde{\Phi}_1) \cdot \frac{\tilde{\nabla}\Theta_0}{\Theta_0}, \quad (\text{C2})$$

where we have used the fact that $\tilde{\nabla}\Theta_0$ has only ξ component and $(\tilde{\Omega} \times \boldsymbol{\xi})_{fig}$ has only φ component. The boundary condition is

$$(\tilde{\nabla}\tilde{\Phi}_1) \cdot (\tilde{\nabla}\Theta_0)|_{\xi_1} = 0. \quad (\text{C3})$$

When we expand $\tilde{\Phi}_1$ as

$$\tilde{\Phi}_1 = \sum_{l,m} {}^{(1)}\phi_{lm}(\xi) \tilde{Y}_l^m(\theta, \varphi), \quad (\text{C4})$$

the boundary conditions are that ${}^{(1)}\phi_{lm}$ is regular at $\xi = 0$ and

$$\frac{d{}^{(1)}\phi_{lm}}{d\xi}(\xi_1) = 0. \quad (\text{C5})$$

We cannot determine the absolute value of $\tilde{\Phi}_1$ from the equation and the boundary condition. Therefore, $\tilde{\Phi}_1$ should be zero. Note that $\tilde{\Phi}_1 = \text{constant}$ is a solution. However, since the physical value is a velocity field $\tilde{\nabla}\tilde{\Phi}_1$, the constant solution is meaningless.

Next, we consider the deformation of the figure at $O(\epsilon)$. The equation is written as

$$\tilde{\Delta}\Theta_1 = -n\Theta_0^{n-1}\Theta_1. \quad (\text{C6})$$

We expand Θ_1 as

$$\Theta_1 = \sum_{l,m} {}^{(1)}\psi_{lm}(\xi) Y_l^m(\theta, \varphi). \quad (\text{C7})$$

The surface equation up to $O(\epsilon)$ is written as

$$\Xi = \xi_1 + \epsilon S_1(\theta, \varphi), \quad (\text{C8})$$

where

$$S_1 = -\frac{\Theta_1(\xi_1)}{\Theta_{0,\xi}(\xi_1)}. \quad (\text{C9})$$

The gravitational potentials and their derivatives up to $O(\epsilon)$ are

$$\tilde{U}_{int} = \Theta - \epsilon \frac{\mu(3+2p)}{2(1+p)^2} + c_0 + \epsilon c_1, \quad (C10)$$

$$\frac{\partial \tilde{U}_{int}}{\partial \xi} = \frac{\partial \Theta}{\partial \xi}, \quad (C11)$$

$$\tilde{U}_{ext} = \frac{\kappa_0}{\xi} + \epsilon \sum_{l,m} \frac{\kappa_{1:lm}}{\xi^{l+1}} Y_l^m, \quad (C12)$$

$$\frac{\partial \tilde{U}_{ext}}{\partial \xi} = -\frac{\kappa_0}{\xi^2} - \epsilon \sum_{l,m} (l+1) \frac{\kappa_{1:lm}}{\xi^{l+2}} Y_l^m. \quad (C13)$$

They become at the stellar surface as

$$\tilde{U}_{int}(\Xi) = c_0 + \epsilon \left[c_1 - \frac{\mu(3+2p)}{2(1+p)^2} \right], \quad (C14)$$

$$\frac{\partial \tilde{U}_{int}}{\partial \xi}(\Xi) = \frac{d\Theta_0}{d\xi}(\xi_1) + \epsilon \left[S_1 \frac{d^2\Theta_0}{d\xi^2}(\xi_1) + \frac{\partial \Theta_1}{\partial \xi}(\xi_1) \right], \quad (C15)$$

$$\tilde{U}_{ext}(\Xi) = \frac{\kappa_0}{\xi_1} \left(1 - \epsilon \frac{S_1}{\xi_1} \right) + \epsilon \sum_{l,m} \frac{\kappa_{1:lm}}{\xi_1^{l+1}} Y_l^m, \quad (C16)$$

$$\frac{\partial \tilde{U}_{ext}}{\partial \xi}(\Xi) = -\frac{\kappa_0}{\xi_1^2} \left(1 - \epsilon \frac{2S_1}{\xi_1} \right) - \epsilon \sum_{l,m} (l+1) \frac{\kappa_{1:lm}}{\xi_1^{l+2}} Y_l^m. \quad (C17)$$

Then, the boundary conditions for $O(\epsilon)$ become

$$c_1 - \frac{\mu(3+2p)}{2(1+p)^2} = -\frac{\kappa_0}{\xi_1^2} S_1 + \sum_{l,m} \frac{\kappa_{1:lm}}{\xi_1^{l+1}} Y_l^m, \quad (C18)$$

$$S_1 \frac{d^2\Theta_0}{d\xi^2}(\xi_1) + \frac{\partial \Theta_1}{\partial \xi}(\xi_1) = \frac{2\kappa_0}{\xi_1^3} S_1 - \sum_{l,m} (l+1) \frac{\kappa_{1:lm}}{\xi_1^{l+2}} Y_l^m, \quad (C19)$$

and also Θ_1 should be satisfied with the conditions at the center of the star,

$$\frac{\partial \Theta_1}{\partial \xi}(0) = 0, \quad (C20)$$

$$\Theta_1(0) = 0. \quad (C21)$$

1. The case of $n = 0$

In this case, since we can express $^{(1)}\psi_{lm} = ^{(1)}A_{lm}\xi^l$ by using a constant $^{(1)}A_{lm}$, the boundary conditions can be written as

$$c_1 - \frac{\mu(3+2p)}{2(1+p)^2} = \sum_{l,m} \left(-^{(1)}A_{lm}\xi_1^l + \frac{\kappa_{1:lm}}{\xi_1^{l+1}} \right) Y_l^m, \quad (C22)$$

$$0 = \sum_{l,m} \left[(3-l)^{(1)}A_{lm}\xi_1^{l-1} - (l+1) \frac{\kappa_{1:lm}}{\xi_1^{l+2}} \right] Y_l^m. \quad (C23)$$

Therefore, we obtain

$$c_1 = \frac{\mu(3+2p)}{2(1+p)^2}, \quad (C24)$$

$$\kappa_{1:lm} = 0, \quad (C25)$$

$$\Theta_1 = 0. \quad (C26)$$

2. The case of $n > 0$

In this case, we have the relation;

$$\frac{d^2\Theta_0}{d\xi^2}(\xi_1) = -\frac{2}{\xi_1} \frac{d\Theta_0}{d\xi}(\xi_1), \quad (\text{C27})$$

from the Lane-Emden equation at the stellar surface. Then, the boundary conditions can be written as

$$c_1 - \frac{\mu(3+2p)}{2(1+p)^2} = \sum_{l,m} \left[-^{(1)}\psi_{lm}(\xi_1) + \frac{\kappa_{1:lm}}{\xi_1^{l+1}} \right] Y_l^m, \quad (\text{C28})$$

$$0 = \sum_{l,m} \left[-\frac{d^{(1)}\psi_{lm}}{d\xi}(\xi_1) - (l+1) \frac{\kappa_{1:lm}}{\xi_1^{l+2}} \right] Y_l^m. \quad (\text{C29})$$

Therefore, we obtain

$$c_1 = \frac{\mu(3+2p)}{2(1+p)^2}, \quad (\text{C30})$$

$$\kappa_{1:lm} = 0, \quad (\text{C31})$$

$$\Theta_1 = 0. \quad (\text{C32})$$

APPENDIX D: SOLUTIONS AT SECOND ORDER

The equation for the velocity potential is

$$\tilde{\Delta}\tilde{\Phi}_2 = -n(\tilde{\nabla}\tilde{\Phi}_2) \cdot \frac{\tilde{\nabla}\Theta_0}{\Theta_0}. \quad (\text{D1})$$

The boundary condition is

$$(\tilde{\nabla}\tilde{\Phi}_2) \cdot (\tilde{\nabla}\Theta_0)|_{\xi_1} = 0. \quad (\text{D2})$$

Then, $\tilde{\Phi}_2$ should be zero as in the case of $\tilde{\Phi}_1$.

Next, we consider the deformation of the figure at $O(\epsilon^2)$. The equation is written as

$$\tilde{\Delta}\Theta_2 = -n\Theta_0^{n-1}\Theta_2. \quad (\text{D3})$$

We expand Θ_2 as

$$\Theta_2 = \sum_{l,m} {}^{(2)}\psi_{lm}(\xi) Y_l^m(\theta, \varphi). \quad (\text{D4})$$

The surface equation up to $O(\epsilon^2)$ is written as

$$\Xi = \xi_1 + \epsilon^2 S_2(\theta, \varphi), \quad (\text{D5})$$

where

$$S_2 = -\frac{\Theta_2(\xi_1)}{\Theta_{0,\xi}(\xi_1)}. \quad (\text{D6})$$

The gravitational potentials and their derivatives up to $O(\epsilon^2)$ are

$$\tilde{U}_{int} = \Theta + c_0 + \epsilon^2 c_2, \quad (D7)$$

$$\frac{\partial \tilde{U}_{int}}{\partial \xi} = \frac{\partial \Theta}{\partial \xi}, \quad (D8)$$

$$\tilde{U}_{ext} = \frac{\kappa_0}{\xi} + \epsilon^2 \sum_{l,m} \frac{\kappa_{2:lm}}{\xi^{l+1}} Y_l^m, \quad (D9)$$

$$\frac{\partial \tilde{U}_{ext}}{\partial \xi} = -\frac{\kappa_0}{\xi^2} - \epsilon^2 \sum_{l,m} (l+1) \frac{\kappa_{2:lm}}{\xi^{l+2}} Y_l^m. \quad (D10)$$

They become at the stellar surface as

$$\tilde{U}_{int}(\Xi) = c_0 + \epsilon^2 c_2, \quad (D11)$$

$$\frac{\partial \tilde{U}_{int}}{\partial \xi}(\Xi) = \frac{d\Theta_0}{d\xi}(\xi_1) + \epsilon^2 \left[S_2 \frac{d^2 \Theta_0}{d\xi^2}(\xi_1) + \frac{\partial \Theta_2}{\partial \xi}(\xi_1) \right], \quad (D12)$$

$$\tilde{U}_{ext}(\Xi) = \frac{\kappa_0}{\xi_1} \left(1 - \epsilon^2 \frac{S_2}{\xi_1} \right) + \epsilon^2 \sum_{l,m} \frac{\kappa_{2:lm}}{\xi_1^{l+1}} Y_l^m, \quad (D13)$$

$$\frac{\partial \tilde{U}_{ext}}{\partial \xi}(\Xi) = -\frac{\kappa_0}{\xi_1^2} \left(1 - \epsilon^2 \frac{2S_2}{\xi_1} \right) - \epsilon^2 \sum_{l,m} (l+1) \frac{\kappa_{2:lm}}{\xi_1^{l+2}} Y_l^m. \quad (D14)$$

Then, the boundary conditions for $O(\epsilon^2)$ become

$$c_2 = -\frac{\kappa_0}{\xi_1^2} S_2 + \sum_{l,m} \frac{\kappa_{2:lm}}{\xi_1^{l+1}} Y_l^m, \quad (D15)$$

$$S_2 \frac{d^2 \Theta_0}{d\xi^2}(\xi_1) + \frac{\partial \Theta_2}{\partial \xi}(\xi_1) = \frac{2\kappa_0}{\xi_1^3} S_2 - \sum_{l,m} (l+1) \frac{\kappa_{2:lm}}{\xi_1^{l+2}} Y_l^m, \quad (D16)$$

and also Θ_2 should be satisfied with the conditions at the center of the star,

$$\frac{\partial \Theta_2}{\partial \xi}(0) = 0, \quad (D17)$$

$$\Theta_2(0) = 0. \quad (D18)$$

1. The case of $n = 0$

In this case, since we can express $^{(2)}\psi_{lm} = ^{(2)}A_{lm}\xi^l$ by using a constant $^{(2)}A_{lm}$, the boundary conditions can be written as

$$c_2 = \sum_{l,m} \left(-^{(2)}A_{lm}\xi_1^l + \frac{\kappa_{2:lm}}{\xi_1^{l+1}} \right) Y_l^m, \quad (D19)$$

$$0 = \sum_{l,m} \left[(3-l)^{(2)}A_{lm}\xi_1^{l-1} - (l+1) \frac{\kappa_{2:lm}}{\xi_1^{l+2}} \right] Y_l^m. \quad (D20)$$

Therefore, we obtain

$$c_2 = 0, \quad (D21)$$

$$\kappa_{2:lm} = 0, \quad (D22)$$

$$\Theta_2 = 0. \quad (D23)$$

2. The case of $n > 0$

In this case, the boundary conditions can be written as

$$c_2 = \sum_{l,m} \left[-^{(2)}\psi_{lm}(\xi_1) + \frac{\kappa_{2:lm}}{\xi_1^{l+1}} \right] Y_l^m, \quad (D24)$$

$$0 = \sum_{l,m} \left[-\frac{d^{(2)}\psi_{lm}}{d\xi}(\xi_1) - (l+1) \frac{\kappa_{2:lm}}{\xi_1^{l+2}} \right] Y_l^m. \quad (D25)$$

Therefore, we obtain

$$c_2 = 0, \quad (D26)$$

$$\kappa_{2:lm} = 0, \quad (D27)$$

$$\Theta_2 = 0. \quad (D28)$$

APPENDIX E: SOLUTIONS AT THIRD ORDER

The equation for the velocity potential is

$$\tilde{\Delta}\tilde{\Phi}_3 = -n(\tilde{\nabla}\tilde{\Phi}_3) \cdot \frac{\tilde{\nabla}\Theta_0}{\Theta_0}. \quad (E1)$$

The boundary condition is

$$(\tilde{\nabla}\tilde{\Phi}_3) \cdot (\tilde{\nabla}\Theta_0)|_{\xi_1} = 0. \quad (E2)$$

Then, $\tilde{\Phi}_3$ should be zero as in the cases of $\tilde{\Phi}_1$ and $\tilde{\Phi}_2$.

Next, we consider the deformation of the figure at $O(\epsilon^3)$. The equation is written as

$$\tilde{\Delta}\Theta_3 = -n\Theta_0^{n-1}\Theta_3. \quad (E3)$$

We expand Θ_3 as

$$\Theta_3 = \sum_{l,m} {}^{(3)}\psi_{lm}(\xi) Y_l^m(\theta, \varphi). \quad (E4)$$

The surface equation up to $O(\epsilon^3)$ is written as

$$\Xi = \xi_1 + \epsilon^3 S_3(\theta, \varphi), \quad (E5)$$

where

$$S_3 = -\frac{\Theta_3(\xi_1)}{\Theta_{0,\xi}(\xi_1)}. \quad (E6)$$

The gravitational potentials and their derivatives up to $O(\epsilon^3)$ are

$$\tilde{U}_{int} = \Theta + c_0 + \epsilon^3 \left[c_3 - \frac{\mu}{\xi_1^2(1+p)} \xi^2 P_2(\sin \theta \cos \varphi) \right], \quad (E7)$$

$$\frac{\partial \tilde{U}_{int}}{\partial \xi} = \frac{\partial \Theta}{\partial \xi} - \epsilon^3 \frac{2\mu}{\xi_1^2(1+p)} \xi P_2(\sin \theta \cos \varphi), \quad (E8)$$

$$\tilde{U}_{ext} = \frac{\kappa_0}{\xi} + \epsilon^3 \sum_{l,m} \frac{\kappa_{3:lm}}{\xi^{l+1}} Y_l^m, \quad (E9)$$

$$\frac{\partial \tilde{U}_{ext}}{\partial \xi} = -\frac{\kappa_0}{\xi^2} - \epsilon^3 \sum_{l,m} (l+1) \frac{\kappa_{3:lm}}{\xi^{l+2}} Y_l^m. \quad (E10)$$

They become at the stellar surface as

$$\tilde{U}_{int}(\Xi) = c_0 + \epsilon^3 \left[c_3 - \frac{\mu}{1+p} P_2(\sin \theta \cos \varphi) \right], \quad (\text{E11})$$

$$\frac{\partial \tilde{U}_{int}}{\partial \xi}(\Xi) = \frac{d\Theta_0}{d\xi}(\xi_1) + \epsilon^3 \left[S_3 \frac{d^2 \Theta_0}{d\xi^2}(\xi_1) + \frac{\partial \Theta_3}{\partial \xi}(\xi_1) - \frac{2\mu}{\xi_1(1+p)} P_2(\sin \theta \cos \varphi) \right], \quad (\text{E12})$$

$$\tilde{U}_{ext}(\Xi) = \frac{\kappa_0}{\xi_1} \left(1 - \epsilon^3 \frac{S_3}{\xi_1} \right) + \epsilon^3 \sum_{l,m} \frac{\kappa_{3:lm}}{\xi_1^{l+1}} Y_l^m, \quad (\text{E13})$$

$$\frac{\partial \tilde{U}_{ext}}{\partial \xi}(\Xi) = -\frac{\kappa_0}{\xi_1^2} \left(1 - \epsilon^3 \frac{2S_3}{\xi_1} \right) - \epsilon^3 \sum_{l,m} (l+1) \frac{\kappa_{3:lm}}{\xi_1^{l+2}} Y_l^m. \quad (\text{E14})$$

Then, the boundary conditions for $O(\epsilon^3)$ become

$$c_3 - \frac{\mu}{1+p} P_2(\sin \theta \cos \varphi) = -\frac{\kappa_0}{\xi_1^2} S_3 + \sum_{l,m} \frac{\kappa_{3:lm}}{\xi_1^{l+1}} Y_l^m, \quad (\text{E15})$$

$$S_3 \frac{d^2 \Theta_0}{d\xi^2}(\xi_1) + \frac{\partial \Theta_3}{\partial \xi}(\xi_1) - \frac{2\mu}{\xi_1(1+p)} P_2(\sin \theta \cos \varphi) = \frac{2\kappa_0}{\xi_1^3} S_3 - \sum_{l,m} (l+1) \frac{\kappa_{3:lm}}{\xi_1^{l+2}} Y_l^m, \quad (\text{E16})$$

and also Θ_3 should be satisfied with the conditions at the center of the star,

$$\frac{\partial \Theta_3}{\partial \xi}(0) = 0, \quad (\text{E17})$$

$$\Theta_3(0) = 0. \quad (\text{E18})$$

1. The case of $n = 0$

In this case, since we can express ${}^{(3)}\psi_{lm} = {}^{(3)}A_{lm}\xi^l$ by using a constant ${}^{(3)}A_{lm}$, the boundary conditions can be written as

$$c_3 - \frac{\mu}{1+p} P_2(\sin \theta \cos \varphi) = \sum_{l,m} \left(-{}^{(3)}A_{lm}\xi_1^l + \frac{\kappa_{3:lm}}{\xi_1^{l+1}} \right) Y_l^m, \quad (\text{E19})$$

$$-\frac{2\mu}{\xi_1(1+p)} P_2(\sin \theta \cos \varphi) = \sum_{l,m} \left[(3-l){}^{(3)}A_{lm}\xi_1^{l-1} - (l+1) \frac{\kappa_{3:lm}}{\xi_1^{l+2}} \right] Y_l^m. \quad (\text{E20})$$

Therefore, we obtain

$$c_3 = 0, \quad (\text{E21})$$

$$\Theta_3 = {}^{(3)}A_2 \xi^2 P_2(\sin \theta \cos \varphi), \quad (\text{E22})$$

where

$${}^{(3)}A_2 = \frac{5\mu}{2(1+p)\xi_1^2}, \quad (\text{E23})$$

$$\kappa_{3:2} = \frac{3\mu\xi_1^3}{2(1+p)}. \quad (\text{E24})$$

2. The case of $n = 1$

In this case, since we can express ${}^{(3)}\psi_{lm} = {}^{(3)}B_{lm}j_l$ by using a constant ${}^{(3)}B_{lm}$ and a spherical Bessel function j_l , the boundary conditions can be written as

$$c_3 - \frac{\mu}{1+p} P_2(\sin \theta \cos \varphi) = \sum_{l,m} \left[-^{(3)}B_{lm} j_l(\xi_1) + \frac{\kappa_{3:lm}}{\xi_1^{l+1}} \right] Y_l^m, \quad (\text{E25})$$

$$-\frac{2\mu}{\xi_1(1+p)} P_2(\sin \theta \cos \varphi) = \sum_{l,m} \left[-^{(3)}B_{lm} \frac{dj_l}{d\xi}(\xi_1) - (l+1) \frac{\kappa_{3:lm}}{\xi_1^{l+2}} \right] Y_l^m. \quad (\text{E26})$$

Therefore, we obtain

$$c_3 = 0, \quad (\text{E27})$$

$$\Theta_3 = ^{(3)}B_2 j_2(\xi) P_2(\sin \theta \cos \varphi), \quad (\text{E28})$$

where

$$^{(3)}B_2 = \frac{5\mu}{1+p}, \quad (\text{E29})$$

$$\kappa_{3:2} = \frac{\mu \xi_1}{1+p} (15 - \xi_1^2), \quad (\text{E30})$$

$$j_2(\xi) = \frac{1}{\xi^3} \left[(3 - \xi^2) \sin \xi - 3\xi \cos \xi \right]. \quad (\text{E31})$$

3. The case of $n > 0$

In this case, the boundary conditions can be written as

$$c_3 - \frac{\mu}{1+p} P_2(\sin \theta \cos \varphi) = \sum_{l,m} \left[-^{(3)}\psi_{lm}(\xi_1) + \frac{\kappa_{3:lm}}{\xi_1^{l+1}} \right] Y_l^m, \quad (\text{E32})$$

$$-\frac{2\mu}{\xi_1(1+p)} P_2(\sin \theta \cos \varphi) = \sum_{l,m} \left[-\frac{d^{(3)}\psi_{lm}}{d\xi}(\xi_1) - (l+1) \frac{\kappa_{3:lm}}{\xi_1^{l+2}} \right] Y_l^m. \quad (\text{E33})$$

Therefore,

$$c_3 = 0, \quad (\text{E34})$$

$$\Theta_3 = ^{(3)}\psi_2(\xi) P_2(\sin \theta \cos \varphi), \quad (\text{E35})$$

and the ordinary differential equation for $^{(3)}\psi_2$ is written as

$$\left[\frac{1}{\xi^2} \frac{d}{d\xi} \left(\xi^2 \frac{d}{d\xi} \right) - \frac{6}{\xi^2} \right] ^{(3)}\psi_2 = -n \Theta_0^{n-1} ^{(3)}\psi_2. \quad (\text{E36})$$

The coefficient of $^{(3)}\psi_2$ and the constant $\kappa_{3:2}$ are determined from two equations;

$$3^{(3)}\psi_2(\xi_1) + \xi_1 \frac{d^{(3)}\psi_2}{d\xi}(\xi_1) = \frac{5\mu}{1+p}, \quad (\text{E37})$$

$$\kappa_{3:2} = \xi_1^3 \left[^{(3)}\psi_2(\xi_1) - \frac{\mu}{1+p} \right] = \frac{\xi_1^3}{3} \left[\frac{2\mu}{1+p} - \xi_1 \frac{d^{(3)}\psi_2}{d\xi}(\xi_1) \right]. \quad (\text{E38})$$

4. Reduced quadrupole moments

At this order, we obtain the reduced quadrupole moments as

$$\mathcal{I}_{11} = \frac{2}{3} \int_{star} d^3 x \rho r^2 P_2(\sin \theta \cos \varphi), \quad (\text{E39})$$

$$= \begin{cases} \epsilon^3 \frac{8\pi}{5} \rho_c \alpha^5 \xi_1^5 A_2, & (n=0) \\ \epsilon^3 \frac{8\pi}{15} \rho_c \alpha^5 \xi_1^3 \left(\frac{15}{\xi_1^2} - 1 \right) ^{(3)}B_2, & (n=1) \\ \epsilon^3 \frac{8\pi}{15} \rho_c \alpha^5 \int_0^{\xi_1} d\xi n \Theta_0^{n-1}(\xi) ^{(3)}\psi_2(\xi) \xi^4. & (n>0) \end{cases} \quad (\text{E40})$$

APPENDIX F: SOLUTIONS AT FOURTH ORDER

The equation for the velocity potential is

$$\tilde{\Delta}\tilde{\Phi}_4 = -\frac{n}{\Theta_0} \left[(\tilde{\nabla}\tilde{\Phi}_4) \cdot (\tilde{\nabla}\Theta_0) - \frac{1}{\xi_1} (\tilde{\mathbf{n}} \times \boldsymbol{\xi})_{fig} \cdot (\tilde{\nabla}\Theta_3) \right], \quad (\text{F1})$$

$$= -\frac{n}{\Theta_0} \left[(\tilde{\nabla}\tilde{\Phi}_4) \cdot (\tilde{\nabla}\Theta_0) + \frac{1}{2\xi_1} {}^{(3)}\psi_2 P_2^2(\cos\theta) \sin 2\varphi \right]. \quad (\text{F2})$$

Therefore, we have the form of $\tilde{\Phi}_4$ and the equation for it as

$$\tilde{\Phi}_4 = {}^{(4)}\phi_2(\xi) P_2^2(\cos\theta) \sin 2\varphi, \quad (\text{F3})$$

$$\left[\frac{1}{\xi^2} \frac{d}{d\xi} \left(\xi^2 \frac{d}{d\xi} \right) - \frac{6}{\xi^2} + \frac{n}{\Theta_0} \frac{d\Theta_0}{d\xi} \frac{d}{d\xi} \right] {}^{(4)}\phi_2 = -\frac{n}{2\xi_1 \Theta_0} {}^{(3)}\psi_2. \quad (\text{F4})$$

The boundary condition is

$$\frac{d^{(4)}\phi_2}{d\xi}(\xi_1) \frac{d\Theta_0}{d\xi}(\xi_1) + \frac{1}{2\xi_1} {}^{(3)}\psi_2(\xi_1) = 0. \quad (\text{F5})$$

Next, we consider the deformation of the figure at $O(\epsilon^4)$. The equation is written as

$$\tilde{\Delta}\Theta_4 = -n\Theta_0^{n-1}\Theta_4. \quad (\text{F6})$$

We expand Θ_4 as

$$\Theta_4 = \sum_{l,m} {}^{(4)}\psi_{lm}(\xi) Y_l^m(\theta, \varphi). \quad (\text{F7})$$

The surface equation up to $O(\epsilon^4)$ is written as

$$\Xi = \xi_1 + \epsilon^3 S_3(\theta, \varphi) + \epsilon^4 S_4(\theta, \varphi), \quad (\text{F8})$$

where

$$S_4 = -\frac{\Theta_4(\xi_1)}{\Theta_{0,\xi}(\xi_1)}. \quad (\text{F9})$$

The gravitational potentials and their derivatives up to $O(\epsilon^4)$ are

$$\tilde{U}_{int} = \Theta + c_0 - \epsilon^3 \frac{\mu}{\xi_1^2(1+p)} \xi^2 P_2(\sin\theta \cos\varphi) + \epsilon^4 \left[c_4 + \frac{\mu}{\xi_1^3(1+p)} \xi^3 P_3(\sin\theta \cos\varphi) \right], \quad (\text{F10})$$

$$\frac{\partial \tilde{U}_{int}}{\partial \xi} = \frac{\partial \Theta}{\partial \xi} - \epsilon^3 \frac{2\mu}{\xi_1^2(1+p)} \xi P_2(\sin\theta \cos\varphi) + \epsilon^4 \frac{3\mu}{\xi_1^3(1+p)} \xi^2 P_3(\sin\theta \cos\varphi), \quad (\text{F11})$$

$$\tilde{U}_{ext} = \frac{\kappa_0}{\xi} + \epsilon^3 \frac{\kappa_{3:2}}{\xi^3} P_2(\sin\theta \cos\varphi) + \epsilon^4 \sum_{l,m} \frac{\kappa_{4:lm}}{\xi^{l+1}} Y_l^m, \quad (\text{F12})$$

$$\frac{\partial \tilde{U}_{ext}}{\partial \xi} = -\frac{\kappa_0}{\xi^2} - \epsilon^3 \frac{3\kappa_{3:2}}{\xi^4} P_2(\sin\theta \cos\varphi) - \epsilon^4 \sum_{l,m} (l+1) \frac{\kappa_{4:lm}}{\xi^{l+2}} Y_l^m. \quad (\text{F13})$$

They become at the stellar surface as

$$\tilde{U}_{int}(\Xi) = c_0 - \epsilon^3 \frac{\mu}{1+p} P_2(\sin\theta \cos\varphi) + \epsilon^4 \left[c_4 + \frac{\mu}{1+p} P_3(\sin\theta \cos\varphi) \right], \quad (\text{F14})$$

$$\begin{aligned} \frac{\partial \tilde{U}_{int}}{\partial \xi}(\Xi) &= \frac{d\Theta_0}{d\xi}(\xi_1) + \epsilon^3 \left[S_3 \frac{d^2 \Theta_0}{d\xi^2}(\xi_1) + \frac{\partial \Theta_3}{\partial \xi}(\xi_1) - \frac{2\mu}{\xi_1(1+p)} P_2(\sin\theta \cos\varphi) \right] \\ &\quad + \epsilon^4 \left[S_4 \frac{d^2 \Theta_0}{d\xi^2}(\xi_1) + \frac{\partial \Theta_4}{\partial \xi}(\xi_1) + \frac{3\mu}{\xi_1(1+p)} P_3(\sin\theta \cos\varphi) \right], \end{aligned} \quad (\text{F15})$$

$$\tilde{U}_{ext}(\Xi) = \frac{\kappa_0}{\xi_1} \left(1 - \epsilon^3 \frac{S_3}{\xi_1} - \epsilon^4 \frac{S_4}{\xi_1} \right) + \epsilon^3 \frac{\kappa_{3:2}}{\xi_1^3} P_2(\sin\theta \cos\varphi) + \epsilon^4 \sum_{l,m} \frac{\kappa_{4:lm}}{\xi_1^{l+1}} Y_l^m, \quad (\text{F16})$$

$$\frac{\partial \tilde{U}_{ext}}{\partial \xi}(\Xi) = -\frac{\kappa_0}{\xi_1^2} \left(1 - \epsilon^3 \frac{2S_3}{\xi_1} - \epsilon^4 \frac{2S_4}{\xi_1} \right) - \epsilon^3 \frac{3\kappa_{3:2}}{\xi_1^4} P_2(\sin\theta \cos\varphi) - \epsilon^4 \sum_{l,m} (l+1) \frac{\kappa_{4:lm}}{\xi_1^{l+2}} Y_l^m. \quad (\text{F17})$$

Then, the boundary conditions for $O(\epsilon^4)$ become

$$c_4 + \frac{\mu}{1+p} P_3(\sin \theta \cos \varphi) = -\frac{\kappa_0}{\xi_1^2} S_4 + \sum_{l,m} \frac{\kappa_{4:lm}}{\xi_1^{l+1}} Y_l^m, \quad (\text{F18})$$

$$S_4 \frac{d^2 \Theta_0}{d\xi^2}(\xi_1) + \frac{\partial \Theta_4}{\partial \xi}(\xi_1) + \frac{3\mu}{\xi_1(1+p)} P_3(\sin \theta \cos \varphi) = \frac{2\kappa_0}{\xi_1^3} S_4 - \sum_{l,m} (l+1) \frac{\kappa_{4:lm}}{\xi_1^{l+2}} Y_l^m, \quad (\text{F19})$$

and also Θ_4 should be satisfied with the conditions at the center of the star,

$$\frac{\partial \Theta_4}{\partial \xi}(0) = 0, \quad (\text{F20})$$

$$\Theta_4(0) = 0. \quad (\text{F21})$$

1. The case of $n = 0$

In this case, since we can express ${}^{(4)}\psi_{lm} = {}^{(4)}A_{lm}\xi^l$ by using a constant ${}^{(4)}A_{lm}$, the boundary conditions can be written as

$$c_4 + \frac{\mu}{1+p} P_3(\sin \theta \cos \varphi) = \sum_{l,m} \left(-{}^{(4)}A_{lm}\xi_1^l + \frac{\kappa_{4:lm}}{\xi_1^{l+1}} \right) Y_l^m, \quad (\text{F22})$$

$$\frac{3\mu}{\xi_1(1+p)} P_3(\sin \theta \cos \varphi) = \sum_{l,m} \left[(3-l){}^{(4)}A_{lm}\xi_1^{l-1} - (l+1)\frac{\kappa_{4:lm}}{\xi_1^{l+2}} \right] Y_l^m. \quad (\text{F23})$$

Therefore, we obtain

$$c_4 = 0, \quad (\text{F24})$$

$$\Theta_4 = {}^{(4)}A_3 \xi^3 P_3(\sin \theta \cos \varphi), \quad (\text{F25})$$

where

$${}^{(4)}A_3 = -\frac{7\mu}{4(1+p)\xi_1^3}, \quad (\text{F26})$$

$$\kappa_{4:3} = -\frac{3\mu\xi_1^4}{4(1+p)}. \quad (\text{F27})$$

2. The case of $n = 1$

In this case, since we can express ${}^{(4)}\psi_{lm} = {}^{(4)}B_{lm}j_l$ by using a constant ${}^{(4)}B_{lm}$, the boundary conditions can be written as

$$c_4 + \frac{\mu}{1+p} P_3(\sin \theta \cos \varphi) = \sum_{l,m} \left[-{}^{(4)}B_{lm}j_l(\xi_1) + \frac{\kappa_{4:lm}}{\xi_1^{l+1}} \right] Y_l^m, \quad (\text{F28})$$

$$\frac{3\mu}{\xi_1(1+p)} P_3(\sin \theta \cos \varphi) = \sum_{l,m} \left[-{}^{(4)}B_{lm} \frac{dj_l}{d\xi}(\xi_1) - (l+1) \frac{\kappa_{4:lm}}{\xi_1^{l+2}} \right] Y_l^m. \quad (\text{F29})$$

Therefore, we obtain

$$c_4 = 0, \quad (\text{F30})$$

$$\Theta_4 = {}^{(4)}B_{3j_3}(\xi) P_3(\sin \theta \cos \varphi), \quad (\text{F31})$$

where

$${}^{(4)}B_3 = -\frac{7\mu\xi_1}{3(1+p)}, \quad (\text{F32})$$

$$\kappa_{4:3} = -\frac{5\mu\xi_1^2}{3(1+p)}(21 - 2\xi_1^2), \quad (\text{F33})$$

$$j_3(\xi) = \frac{1}{\xi^4} \left[(15 - 6\xi^2) \sin \xi - \xi(15 - \xi^2) \cos \xi \right]. \quad (\text{F34})$$

3. The case of $n > 0$

In this case, the boundary conditions can be written as

$$c_4 + \frac{\mu}{1+p} P_3(\sin \theta \cos \varphi) = \sum_{l,m} \left[-{}^{(4)}\psi_{lm}(\xi_1) + \frac{\kappa_{4:lm}}{\xi_1^{l+1}} \right] Y_l^m, \quad (\text{F35})$$

$$\frac{3\mu}{\xi_1(1+p)} P_3(\sin \theta \cos \varphi) = \sum_{l,m} \left[-\frac{d{}^{(4)}\psi_{lm}}{d\xi}(\xi_1) - (l+1) \frac{\kappa_{4:lm}}{\xi_1^{l+2}} \right] Y_l^m. \quad (\text{F36})$$

Therefore, we obtain

$$c_4 = 0, \quad (\text{F37})$$

$$\Theta_4 = {}^{(4)}\psi_3(\xi) P_3(\sin \theta \cos \varphi), \quad (\text{F38})$$

and the ordinary differential equation for ${}^{(4)}\psi_3$ is written as

$$\left[\frac{1}{\xi^2} \frac{d}{d\xi} \left(\xi^2 \frac{d}{d\xi} \right) - \frac{12}{\xi^2} \right] {}^{(4)}\psi_3 = -n\Theta_0^{n-1} {}^{(4)}\psi_3. \quad (\text{F39})$$

The coefficient of ${}^{(4)}\psi_3$ and the constant $\kappa_{4:3}$ are determined from two equations;

$$4{}^{(4)}\psi_3(\xi_1) + \xi_1 \frac{d{}^{(4)}\psi_3}{d\xi}(\xi_1) = -\frac{7\mu}{1+p}, \quad (\text{F40})$$

$$\kappa_{4:3} = \xi_1^4 \left[{}^{(4)}\psi_3(\xi_1) + \frac{\mu}{1+p} \right] = -\frac{\xi_1^4}{4} \left[\frac{3\mu}{1+p} + \xi_1 \frac{d{}^{(4)}\psi_3}{d\xi}(\xi_1) \right]. \quad (\text{F41})$$

4. Octupole moments

At this order, we can obtain the octupole moments, but we do not represent them here.

APPENDIX G: SOLUTIONS AT FIFTH ORDER

The equation for the velocity potential is

$$\tilde{\Delta} \tilde{\Phi}_5 = -\frac{n}{\Theta_0} \left[(\tilde{\nabla} \tilde{\Phi}_5) \cdot (\tilde{\nabla} \Theta_0) - \frac{1}{\xi_1} (\tilde{\mathbf{\Omega}} \times \boldsymbol{\xi})_{fig} \cdot (\tilde{\nabla} \Theta_4) \right], \quad (\text{G1})$$

$$= -\frac{n}{\Theta_0} \left[(\tilde{\nabla} \tilde{\Phi}_5) \cdot (\tilde{\nabla} \Theta_0) - \frac{1}{4\xi_1} {}^{(4)}\psi_3 \left\{ P_3^1(\cos \theta) \sin \varphi - \frac{1}{2} P_3^3(\cos \theta) \sin 3\varphi \right\} \right]. \quad (\text{G2})$$

Therefore, we have the form of $\tilde{\Phi}_5$ and the equation for it as

$$\tilde{\Phi}_5 = {}^{(5)}\phi_3(\xi) \left[P_3^1(\cos \theta) \sin \varphi - \frac{1}{2} P_3^3(\cos \theta) \sin 3\varphi \right], \quad (\text{G3})$$

$$\left[\frac{1}{\xi^2} \frac{d}{d\xi} \left(\xi^2 \frac{d}{d\xi} \right) - \frac{12}{\xi^2} + \frac{n}{\Theta_0} \frac{d\Theta_0}{d\xi} \frac{d}{d\xi} \right] {}^{(5)}\phi_3 = \frac{n}{4\xi_1 \Theta_0} {}^{(4)}\psi_3. \quad (\text{G4})$$

The boundary condition is

$$\frac{d^{(5)}\phi_3}{d\xi}(\xi_1)\frac{d\Theta_0}{d\xi}(\xi_1) - \frac{1}{4\xi_1}{}^{(4)}\psi_3(\xi_1) = 0. \quad (\text{G5})$$

Next, we consider the deformation of the figure at $O(\epsilon^5)$. The equation is written as

$$\tilde{\Delta}\Theta_5 = -n\Theta_0^{n-1}\Theta_5. \quad (\text{G6})$$

We expand Θ_5 as

$$\Theta_5 = \sum_{l,m} {}^{(5)}\psi_{lm}(\xi)Y_l^m(\theta, \varphi). \quad (\text{G7})$$

The surface equation up to $O(\epsilon^5)$ is written as

$$\Xi = \xi_1 + \epsilon^3 S_3(\theta, \varphi) + \epsilon^4 S_4(\theta, \varphi) + \epsilon^5 S_5(\theta, \varphi), \quad (\text{G8})$$

where

$$S_5 = -\frac{\Theta_5(\xi_1)}{\Theta_{0,\xi}(\xi_1)}. \quad (\text{G9})$$

The gravitational potentials and their derivatives up to $O(\epsilon^5)$ are

$$\begin{aligned} \tilde{U}_{int} = & \Theta + c_0 - \epsilon^3 \frac{\mu}{\xi_1^2(1+p)} \xi^2 P_2(\sin \theta \cos \varphi) + \epsilon^4 \frac{\mu}{\xi_1^3(1+p)} \xi^3 P_3(\sin \theta \cos \varphi) \\ & + \epsilon^5 \left[c_5 - \frac{\mu}{\xi_1^4(1+p)} \xi^4 P_4(\sin \theta \cos \varphi) \right], \end{aligned} \quad (\text{G10})$$

$$\begin{aligned} \frac{\partial \tilde{U}_{int}}{\partial \xi} = & \frac{\partial \Theta}{\partial \xi} - \epsilon^3 \frac{2\mu}{\xi_1^2(1+p)} \xi P_2(\sin \theta \cos \varphi) + \epsilon^4 \frac{3\mu}{\xi_1^3(1+p)} \xi^2 P_3(\sin \theta \cos \varphi) \\ & - \epsilon^5 \frac{4\mu}{\xi_1^4(1+p)} \xi^3 P_4(\sin \theta \cos \varphi), \end{aligned} \quad (\text{G11})$$

$$\tilde{U}_{ext} = \frac{\kappa_0}{\xi} + \epsilon^3 \frac{\kappa_{3:2}}{\xi^3} P_2(\sin \theta \cos \varphi) + \epsilon^4 \frac{\kappa_{4:3}}{\xi^4} P_3(\sin \theta \cos \varphi) + \epsilon^5 \sum_{l,m} \frac{\kappa_{5:lm}}{\xi^{l+1}} Y_l^m, \quad (\text{G12})$$

$$\frac{\partial \tilde{U}_{ext}}{\partial \xi} = -\frac{\kappa_0}{\xi^2} - \epsilon^3 \frac{3\kappa_{3:2}}{\xi^4} P_2(\sin \theta \cos \varphi) - \epsilon^4 \frac{4\kappa_{4:3}}{\xi^5} P_3(\sin \theta \cos \varphi) - \epsilon^5 \sum_{l,m} (l+1) \frac{\kappa_{5:lm}}{\xi^{l+2}} Y_l^m. \quad (\text{G13})$$

They become at the stellar surface as

$$\tilde{U}_{int}(\Xi) = c_0 - \epsilon^3 \frac{\mu}{1+p} P_2(\sin \theta \cos \varphi) + \epsilon^4 \frac{\mu}{1+p} P_3(\sin \theta \cos \varphi) + \epsilon^5 \left[c_5 - \frac{\mu}{1+p} P_4(\sin \theta \cos \varphi) \right], \quad (\text{G14})$$

$$\begin{aligned} \frac{\partial \tilde{U}_{int}}{\partial \xi}(\Xi) = & \frac{d\Theta_0}{d\xi}(\xi_1) + \epsilon^3 \left[S_3 \frac{d^2\Theta_0}{d\xi^2}(\xi_1) + \frac{\partial\Theta_3}{\partial\xi}(\xi_1) - \frac{2\mu}{\xi_1(1+p)} P_2(\sin \theta \cos \varphi) \right] \\ & + \epsilon^4 \left[S_4 \frac{d^2\Theta_0}{d\xi^2}(\xi_1) + \frac{\partial\Theta_4}{\partial\xi}(\xi_1) + \frac{3\mu}{\xi_1(1+p)} P_3(\sin \theta \cos \varphi) \right] \\ & + \epsilon^5 \left[S_5 \frac{d^2\Theta_0}{d\xi^2}(\xi_1) + \frac{\partial\Theta_5}{\partial\xi}(\xi_1) - \frac{4\mu}{\xi_1(1+p)} P_4(\sin \theta \cos \varphi) \right], \end{aligned} \quad (\text{G15})$$

$$\tilde{U}_{ext}(\Xi) = \frac{\kappa_0}{\xi_1} \left(1 - \epsilon^3 \frac{S_3}{\xi_1} - \epsilon^4 \frac{S_4}{\xi_1} - \epsilon^5 \frac{S_5}{\xi_1} \right) + \epsilon^3 \frac{\kappa_{3:2}}{\xi_1^3} P_2(\sin \theta \cos \varphi) + \epsilon^4 \frac{\kappa_{4:3}}{\xi_1^4} P_3(\sin \theta \cos \varphi) + \epsilon^5 \sum_{l,m} \frac{\kappa_{5:lm}}{\xi_1^{l+1}} Y_l^m, \quad (\text{G16})$$

$$\begin{aligned} \frac{\partial \tilde{U}_{ext}}{\partial \xi}(\Xi) = & -\frac{\kappa_0}{\xi_1^2} \left(1 - \epsilon^3 \frac{2S_3}{\xi_1} - \epsilon^4 \frac{2S_4}{\xi_1} - \epsilon^5 \frac{2S_5}{\xi_1} \right) - \epsilon^3 \frac{3\kappa_{3:2}}{\xi_1^4} P_2(\sin \theta \cos \varphi) - \epsilon^4 \frac{4\kappa_{4:3}}{\xi_1^5} P_3(\sin \theta \cos \varphi) \\ & - \epsilon^5 \sum_{l,m} (l+1) \frac{\kappa_{5:lm}}{\xi_1^{l+2}} Y_l^m. \end{aligned} \quad (\text{G17})$$

Then, the boundary conditions for $O(\epsilon^5)$ become

$$c_5 - \frac{\mu}{1+p} P_4(\sin \theta \cos \varphi) = -\frac{\kappa_0}{\xi_1^2} S_5 + \sum_{l,m} \frac{\kappa_{5:lm}}{\xi_1^{l+1}} Y_l^m, \quad (\text{G18})$$

$$S_5 \frac{d^2 \Theta_0}{d\xi^2}(\xi_1) + \frac{\partial \Theta_5}{\partial \xi}(\xi_1) - \frac{4\mu}{\xi_1(1+p)} P_4(\sin \theta \cos \varphi) = \frac{2\kappa_0}{\xi_1^3} S_5 - \sum_{l,m} (l+1) \frac{\kappa_{5:lm}}{\xi_1^{l+2}} Y_l^m, \quad (\text{G19})$$

and also Θ_5 should be satisfied with the conditions at the center of the star,

$$\frac{\partial \Theta_5}{\partial \xi}(0) = 0, \quad (\text{G20})$$

$$\Theta_5(0) = 0. \quad (\text{G21})$$

1. The case of $n = 0$

In this case, since we can express ${}^{(5)}\psi_{lm} = {}^{(5)}A_{lm}\xi^l$ by using a constant ${}^{(5)}A_{lm}$, the boundary conditions can be written as

$$c_5 - \frac{\mu}{1+p} P_4(\sin \theta \cos \varphi) = \sum_{l,m} \left(-{}^{(5)}A_{lm}\xi_1^l + \frac{\kappa_{5:lm}}{\xi_1^{l+1}} \right) Y_l^m, \quad (\text{G22})$$

$$-\frac{4\mu}{\xi_1(1+p)} P_4(\sin \theta \cos \varphi) = \sum_{l,m} \left[(3-l){}^{(5)}A_{lm}\xi_1^{l-1} - (l+1) \frac{\kappa_{5:lm}}{\xi_1^{l+2}} \right] Y_l^m. \quad (\text{G23})$$

Therefore, we obtain

$$c_5 = 0, \quad (\text{G24})$$

$$\Theta_5 = {}^{(5)}A_4 \xi^4 P_4(\sin \theta \cos \varphi), \quad (\text{G25})$$

where

$${}^{(5)}A_4 = \frac{3\mu}{2(1+p)\xi_1^4}, \quad (\text{G26})$$

$$\kappa_{5:4} = \frac{\mu\xi_1^5}{2(1+p)}. \quad (\text{G27})$$

2. The case of $n = 1$

In this case, since we can express ${}^{(5)}\psi_{lm} = {}^{(5)}B_{lm}j_l$ by using a constant ${}^{(5)}B_{lm}$, the boundary conditions can be written as

$$c_5 - \frac{\mu}{1+p} P_4(\sin \theta \cos \varphi) = \sum_{l,m} \left[-{}^{(5)}B_{lm}j_l(\xi_1) + \frac{\kappa_{5:lm}}{\xi_1^{l+1}} \right] Y_l^m, \quad (\text{G28})$$

$$-\frac{4\mu}{\xi_1(1+p)} P_4(\sin \theta \cos \varphi) = \sum_{l,m} \left[-{}^{(5)}B_{lm} \frac{dj_l}{d\xi}(\xi_1) - (l+1) \frac{\kappa_{5:lm}}{\xi_1^{l+2}} \right] Y_l^m. \quad (\text{G29})$$

Therefore, we obtain

$$c_5 = 0, \quad (\text{G30})$$

$$\Theta_5 = {}^{(5)}B_4 j_4(\xi) P_4(\sin \theta \cos \varphi), \quad (\text{G31})$$

where

$${}^{(5)}B_4 = \frac{9\mu\xi_1^2}{(1+p)(15-\xi_1^2)}, \quad (G32)$$

$$\kappa_{5:4} = \frac{\mu}{1+p} \left(\frac{\xi_1^4 - 105\xi_1^2 + 945}{\xi_1^2(15-\xi_1^2)} \right), \quad (G33)$$

$$j_4(\xi) = \frac{1}{\xi^5} \left[(105 - 45\xi^2 + \xi^4) \sin \xi - \xi(105 - 10\xi^2) \cos \xi \right]. \quad (G34)$$

3. The case of $n > 0$

In this case, the boundary conditions can be written as

$$c_5 - \frac{\mu}{1+p} P_4(\sin \theta \cos \varphi) = \sum_{l,m} \left[-{}^{(5)}\psi_{lm}(\xi_1) + \frac{\kappa_{5:lm}}{\xi_1^{l+1}} \right] Y_l^m, \quad (G35)$$

$$-\frac{4\mu}{\xi_1(1+p)} P_4(\sin \theta \cos \varphi) = \sum_{l,m} \left[-\frac{d^{(5)}\psi_{lm}}{d\xi}(\xi_1) - (l+1) \frac{\kappa_{5:lm}}{\xi_1^{l+2}} \right] Y_l^m. \quad (G36)$$

Therefore, we obtain

$$c_5 = 0, \quad (G37)$$

$$\Theta_5 = {}^{(5)}\psi_4(\xi) P_4(\sin \theta \cos \varphi), \quad (G38)$$

and the ordinary differential equation for ${}^{(5)}\psi_4$ is written as

$$\left[\frac{1}{\xi^2} \frac{d}{d\xi} \left(\xi^2 \frac{d}{d\xi} \right) - \frac{20}{\xi^2} \right] {}^{(5)}\psi_4 = -n\Theta_0^{n-1} {}^{(5)}\psi_4. \quad (G39)$$

The coefficient of ${}^{(5)}\psi_4$ and the constant $\kappa_{5:4}$ are determined from two equations;

$$5{}^{(5)}\psi_4(\xi_1) + \xi_1 \frac{d^{(5)}\psi_4}{d\xi}(\xi_1) = \frac{9\mu}{1+p}, \quad (G40)$$

$$\kappa_{5:4} = \xi_1^5 \left[{}^{(5)}\psi_4(\xi_1) - \frac{\mu}{1+p} \right] = \frac{\xi_1^5}{5} \left[\frac{4\mu}{1+p} - \xi_1 \frac{d^{(5)}\psi_4}{d\xi}(\xi_1) \right]. \quad (G41)$$

4. Orbital angular velocity

At this order, the orbital angular velocity has the 5th higher order term than the monopole term. The total angular velocity is

$$\Omega^2 = \frac{M_{tot}}{R^3} \left[1 + \frac{9}{2R^2} \left(\epsilon^3 \frac{\bar{F}_{11}}{M_1} + \epsilon'^3 \frac{\bar{F}'_{11}}{M_2} \right) \right]. \quad (G42)$$

For the calculational convenience, we rewrite the above equation as

$$\Omega^2 = \frac{M_{tot}}{R^3} (1 + \epsilon^5 \delta), \quad (G43)$$

where

$$\delta \equiv \frac{9}{2a_0^2} \left[\frac{\bar{F}_{11}}{M_1} + \left(\frac{a'_0}{a_0} \right)^3 \frac{\bar{F}'_{11}}{M_2} \right]. \quad (G44)$$

APPENDIX H: SOLUTIONS AT SIXTH ORDER

The equation for the velocity potential is

$$\tilde{\Delta}\tilde{\Phi}_6 = -\frac{n}{\Theta_0} \left[(\tilde{\nabla}\tilde{\Phi}_6) \cdot (\tilde{\nabla}\Theta_0) - \frac{1}{\xi_1} (\tilde{\Omega} \times \xi)_{fig} \cdot (\tilde{\nabla}\Theta_5) \right], \quad (H1)$$

$$= -\frac{n}{\Theta_0} \left[(\tilde{\nabla}\tilde{\Phi}_6) \cdot (\tilde{\nabla}\Theta_0) - \frac{1}{12\xi_1} {}^{(5)}\psi_4 \left\{ P_4^2(\cos\theta) \sin 2\varphi - \frac{1}{4} P_4^4(\cos\theta) \sin 4\varphi \right\} \right]. \quad (H2)$$

Therefore, we have the form of $\tilde{\Phi}_6$ and the equation for it as

$$\tilde{\Phi}_6 = {}^{(6)}\phi_4(\xi) \left[P_4^2(\cos\theta) \sin 2\varphi - \frac{1}{4} P_4^4(\cos\theta) \sin 4\varphi \right], \quad (H3)$$

$$\left[\frac{1}{\xi^2} \frac{d}{d\xi} \left(\xi^2 \frac{d}{d\xi} \right) - \frac{20}{\xi^2} + \frac{n}{\Theta_0} \frac{d\Theta_0}{d\xi} \frac{d}{d\xi} \right] {}^{(6)}\phi_4 = \frac{n}{12\xi_1\Theta_0} {}^{(5)}\psi_4. \quad (H4)$$

The boundary condition is

$$\frac{d({}^{(6)}\phi_4)}{d\xi}(\xi_1) \frac{d\Theta_0}{d\xi}(\xi_1) - \frac{1}{12\xi_1} {}^{(5)}\psi_4(\xi_1) = 0. \quad (H5)$$

Next, we consider the deformation of the figure at $O(\epsilon^6)$. The equation is written as

$$\tilde{\Delta}\Theta_6 = -n\Theta_0^{n-1}\Theta_6 - \frac{1}{2}n(n-1)\Theta_0^{n-2}\Theta_3^2 + \frac{2\mu}{\xi_1} \tilde{\Delta} {}^{(4)}\phi_2(\xi) P_2^2(\cos\theta) \cos 2\varphi. \quad (H6)$$

We expand Θ_6 as

$$\Theta_6 = \sum_{l,m} {}^{(6)}\psi_{lm}(\xi) Y_l^m(\theta, \varphi). \quad (H7)$$

The surface equation up to $O(\epsilon^6)$ is written as

$$\Xi = \xi_1 + \epsilon^3 S_3(\theta, \varphi) + \epsilon^4 S_4(\theta, \varphi) + \epsilon^5 S_5(\theta, \varphi) + \epsilon^6 S_6(\theta, \varphi), \quad (H8)$$

where

$$S_6 = -\frac{1}{\Theta_{0,\xi}(\xi_1)} \left[\frac{S_3^2}{2} \frac{d^2\Theta_0}{d\xi^2}(\xi_1) + S_3 \frac{\partial\Theta_3}{\partial\xi}(\xi_1) + \Theta_6(\xi_1) \right]. \quad (H9)$$

The gravitational potentials and their derivatives up to $O(\epsilon^6)$ are

$$\begin{aligned} \tilde{U}_{int} &= \Theta + c_0 - \epsilon^3 \frac{\mu}{\xi_1^2(1+p)} \xi^2 P_2(\sin\theta \cos\varphi) + \epsilon^4 \frac{\mu}{\xi_1^3(1+p)} \xi^3 P_3(\sin\theta \cos\varphi) \\ &\quad - \epsilon^5 \frac{\mu}{\xi_1^4(1+p)} \xi^4 P_4(\sin\theta \cos\varphi) \\ &\quad + \epsilon^6 \left[c_6 - \frac{3\mu \bar{F}'_{11}}{2M_{tot}a_0^2} \left(\frac{a'_0}{a_0} \right)^3 - \frac{\mu\delta}{2(1+p)^2} + \frac{\mu}{\xi_1^5(1+p)} \xi^5 P_5(\sin\theta \cos\varphi) - \frac{2\mu}{\xi_1} {}^{(4)}\phi_2 P_2^2(\cos\theta) \cos 2\varphi \right], \end{aligned} \quad (H10)$$

$$\begin{aligned} \frac{\partial\tilde{U}_{int}}{\partial\xi} &= \frac{\partial\Theta}{\partial\xi} - \epsilon^3 \frac{2\mu}{\xi_1^2(1+p)} \xi P_2(\sin\theta \cos\varphi) + \epsilon^4 \frac{3\mu}{\xi_1^3(1+p)} \xi^2 P_3(\sin\theta \cos\varphi) - \epsilon^5 \frac{4\mu}{\xi_1^4(1+p)} \xi^3 P_4(\sin\theta \cos\varphi) \\ &\quad + \epsilon^6 \left[\frac{5\mu}{\xi_1^5(1+p)} \xi^4 P_5(\sin\theta \cos\varphi) - \frac{2\mu}{\xi_1} \frac{d^{(4)}\phi_2}{d\xi} P_2^2(\cos\theta) \cos 2\varphi \right], \end{aligned} \quad (H11)$$

$$\tilde{U}_{ext} = \frac{\kappa_0}{\xi} + \epsilon^3 \frac{\kappa_{3:2}}{\xi^3} P_2(\sin\theta \cos\varphi) + \epsilon^4 \frac{\kappa_{4:3}}{\xi^4} P_3(\sin\theta \cos\varphi) + \epsilon^5 \frac{\kappa_{5:4}}{\xi^5} P_4(\sin\theta \cos\varphi) + \epsilon^6 \sum_{l,m} \frac{\kappa_{6:lm}}{\xi^{l+1}} Y_l^m, \quad (H12)$$

$$\begin{aligned} \frac{\partial\tilde{U}_{ext}}{\partial\xi} &= -\frac{\kappa_0}{\xi^2} - \epsilon^3 \frac{3\kappa_{3:2}}{\xi^4} P_2(\sin\theta \cos\varphi) - \epsilon^4 \frac{4\kappa_{4:3}}{\xi^5} P_3(\sin\theta \cos\varphi) - \epsilon^5 \frac{5\kappa_{5:4}}{\xi^6} P_4(\sin\theta \cos\varphi) \\ &\quad - \epsilon^6 \sum_{l,m} (l+1) \frac{\kappa_{6:lm}}{\xi^{l+2}} Y_l^m. \end{aligned} \quad (H13)$$

They become at the stellar surface as

$$\begin{aligned}\tilde{U}_{int}(\Xi) = & c_0 - \epsilon^3 \frac{\mu}{1+p} \left(1 + \frac{2S_3}{\xi_1} \epsilon^3\right) P_2(\sin \theta \cos \varphi) + \epsilon^4 \frac{\mu}{1+p} P_3(\sin \theta \cos \varphi) - \epsilon^5 \frac{\mu}{1+p} P_4(\sin \theta \cos \varphi) \\ & + \epsilon^6 \left[c_6 - \frac{3\mu \bar{I}'_{11}}{2M_{tot}a_0^2} \left(\frac{a'_0}{a_0}\right)^3 - \frac{\mu\delta}{2(1+p)^2} + \frac{\mu}{1+p} P_5(\sin \theta \cos \varphi) - \frac{2\mu^{(4)}\phi_2(\xi_1)}{\xi_1} P_2^2(\cos \theta) \cos 2\varphi \right],\end{aligned}\quad (H14)$$

$$\begin{aligned}\frac{\partial \tilde{U}_{int}}{\partial \xi}(\Xi) = & \frac{d\Theta_0}{d\xi}(\xi_1) + \epsilon^3 \left[S_3 \frac{d^2\Theta_0}{d\xi^2}(\xi_1) + \frac{\partial\Theta_3}{\partial\xi}(\xi_1) - \frac{2\mu}{\xi_1(1+p)} \left(1 + \frac{S_3}{\xi_1} \epsilon^3\right) P_2(\sin \theta \cos \varphi) \right] \\ & + \epsilon^4 \left[S_4 \frac{d^2\Theta_0}{d\xi^2}(\xi_1) + \frac{\partial\Theta_4}{\partial\xi}(\xi_1) + \frac{3\mu}{\xi_1(1+p)} P_3(\sin \theta \cos \varphi) \right] \\ & + \epsilon^5 \left[S_5 \frac{d^2\Theta_0}{d\xi^2}(\xi_1) + \frac{\partial\Theta_5}{\partial\xi}(\xi_1) - \frac{4\mu}{\xi_1(1+p)} P_4(\sin \theta \cos \varphi) \right] \\ & + \epsilon^6 \left[S_6 \frac{d^2\Theta_0}{d\xi^2}(\xi_1) + \frac{1}{2} S_3^2 \frac{d^3\Theta_0}{d\xi^3}(\xi_1) + S_3 \frac{\partial^2\Theta_3}{\partial\xi^2}(\xi_1) + \frac{\partial\Theta_6}{\partial\xi}(\xi_1) + \frac{5\mu}{\xi_1(1+p)} P_5(\sin \theta \cos \varphi) \right. \\ & \left. - \frac{2\mu}{\xi_1} \frac{d^{(4)}\phi_2}{d\xi}(\xi_1) P_2^2(\cos \theta) \cos 2\varphi \right],\end{aligned}\quad (H15)$$

$$\begin{aligned}\tilde{U}_{ext}(\Xi) = & \frac{\kappa_0}{\xi_1} \left[1 - \epsilon^3 \frac{S_3}{\xi_1} - \epsilon^4 \frac{S_4}{\xi_1} - \epsilon^5 \frac{S_5}{\xi_1} + \epsilon^6 \left\{ \left(\frac{S_3}{\xi_1} \right)^2 - \frac{S_6}{\xi_1} \right\} \right] + \epsilon^3 \left(1 - \epsilon^3 \frac{3S_3}{\xi_1} \right) \frac{\kappa_{3:2}}{\xi_1^3} P_2(\sin \theta \cos \varphi) \\ & + \epsilon^4 \frac{\kappa_{4:3}}{\xi_1^4} P_3(\sin \theta \cos \varphi) + \epsilon^5 \frac{\kappa_{5:4}}{\xi_1^5} P_4(\sin \theta \cos \varphi) + \epsilon^6 \sum_{l,m} \frac{\kappa_{6:lm}}{\xi_1^{l+1}} Y_l^m,\end{aligned}\quad (H16)$$

$$\begin{aligned}\frac{\partial \tilde{U}_{ext}}{\partial \xi}(\Xi) = & -\frac{\kappa_0}{\xi_1^2} \left[1 - \epsilon^3 \frac{2S_3}{\xi_1} - \epsilon^4 \frac{2S_4}{\xi_1} - \epsilon^5 \frac{2S_5}{\xi_1} + \epsilon^6 \left\{ 3 \left(\frac{S_3}{\xi_1} \right)^2 - 2 \frac{S_6}{\xi_1} \right\} \right] - \epsilon^3 \left(1 - \epsilon^3 \frac{4S_3}{\xi_1} \right) \frac{3\kappa_{3:2}}{\xi_1^4} P_2(\sin \theta \cos \varphi) \\ & - \epsilon^4 \frac{4\kappa_{4:3}}{\xi_1^5} P_3(\sin \theta \cos \varphi) - \epsilon^5 \frac{5\kappa_{5:4}}{\xi_1^6} P_4(\sin \theta \cos \varphi) - \epsilon^6 \sum_{l,m} (l+1) \frac{\kappa_{6:lm}}{\xi_1^{l+2}} Y_l^m.\end{aligned}\quad (H17)$$

Then, the boundary conditions for $O(\epsilon^6)$ become

$$\begin{aligned}\tilde{c}_6 - \frac{2\mu}{\xi_1(1+p)} S_3 P_2(\sin \theta \cos \varphi) + \frac{\mu}{1+p} P_5(\sin \theta \cos \varphi) - \frac{2\mu^{(4)}\phi_2(\xi_1)}{\xi_1} P_2^2(\cos \theta) \cos 2\varphi \\ = \frac{\kappa_0}{\xi_1^2} \left(\frac{S_3^2}{\xi_1} - S_6 \right) - \frac{3\kappa_{3:2}}{\xi_1^4} S_3 P_2(\sin \theta \cos \varphi) + \sum_{l,m} \frac{\kappa_{6:lm}}{\xi_1^{l+1}} Y_l^m,\end{aligned}\quad (H18)$$

$$\begin{aligned}S_6 \frac{d^2\Theta_0}{d\xi^2}(\xi_1) + \frac{1}{2} S_3^2 \frac{d^3\Theta_0}{d\xi^3}(\xi_1) + S_3 \frac{\partial^2\Theta_3}{\partial\xi^2}(\xi_1) + \frac{\partial\Theta_6}{\partial\xi}(\xi_1) - \frac{2\mu}{\xi_1^2(1+p)} S_3 P_2(\sin \theta \cos \varphi) + \frac{5\mu}{\xi_1(1+p)} P_5(\sin \theta \cos \varphi) \\ - \frac{2\mu}{\xi_1} \frac{d^{(4)}\phi_2}{d\xi}(\xi_1) P_2^2(\cos \theta) \cos 2\varphi = -\frac{\kappa_0}{\xi_1^3} \left(\frac{3S_3^2}{\xi_1} - 2S_6 \right) + \frac{12\kappa_{3:2}}{\xi_1^5} S_3 P_2(\sin \theta \cos \varphi) - \sum_{l,m} (l+1) \frac{\kappa_{6:lm}}{\xi_1^{l+2}} Y_l^m,\end{aligned}\quad (H19)$$

where

$$\tilde{c}_6 \equiv c_6 - \frac{3\mu \bar{I}'_{11}}{2M_{tot}a_0^2} \left(\frac{a'_0}{a_0}\right)^3 - \frac{\mu\delta}{2(1+p)^2}.\quad (H20)$$

Also Θ_6 should be satisfied with the conditions at the center of the star,

$$\frac{\partial\Theta_6}{\partial\xi}(0) = 0,\quad (H21)$$

$$\Theta_6(0) = 0.\quad (H22)$$

In the calculation of this section, we need the expression of $[P_2(\sin \theta \cos \varphi)]^2$. It is written as

$$[P_2(\sin \theta \cos \varphi)]^2 = \frac{18}{35} P_4(\sin \theta \cos \varphi) + \frac{2}{7} P_2(\sin \theta \cos \varphi) + \frac{1}{5}.\quad (H23)$$

1. The case of $n = 0$

In this case, since we can express ${}^{(3)}\psi_2 = {}^{(3)}A_2\xi^2$ and ${}^{(6)}\psi_{lm} = {}^{(6)}A_{lm}\xi^l$, by using the constants ${}^{(3)}A_2$ and ${}^{(6)}A_{lm}$, the boundary conditions can be written as

$$\begin{aligned} \tilde{c}_6 + \frac{\mu}{1+p}P_5(\sin\theta\cos\varphi) - \frac{2\mu}{\xi_1}{}^{(4)}\phi_2(\xi_1)P_2^2(\cos\theta)\cos 2\varphi \\ = \frac{{}^{(3)}A_2\xi_1^2}{\Theta_{0,\xi}(\xi_1)}\left[-\frac{2\mu}{\xi_1(1+p)} + \frac{1}{2}{}^{(3)}A_2\xi_1 + \frac{3\kappa_{3:2}}{\xi_1^4}\right]\left(\frac{18}{35}P_4(\sin\theta\cos\varphi) + \frac{2}{7}P_2(\sin\theta\cos\varphi) + \frac{1}{5}\right) \\ + \sum_{l,m}\left(-{}^{(6)}A_{lm}\xi_1^l + \frac{\kappa_{6:lm}}{\xi_1^{l+1}}\right)Y_l^m, \end{aligned} \quad (\text{H24})$$

$$\begin{aligned} \frac{5\mu}{\xi_1(1+p)}P_5(\sin\theta\cos\varphi) - \frac{2\mu}{\xi_1}\frac{d^{(4)}\phi_2}{d\xi}(\xi_1)P_2^2(\cos\theta)\cos 2\varphi \\ = \frac{{}^{(3)}A_2\xi_1^2}{\Theta_{0,\xi}(\xi_1)}\left[-\frac{2\mu}{\xi_1^2(1+p)} + \frac{1}{2}{}^{(3)}A_2 - \frac{12\kappa_{3:2}}{\xi_1^5}\right]\left(\frac{18}{35}P_4(\sin\theta\cos\varphi) + \frac{2}{7}P_2(\sin\theta\cos\varphi) + \frac{1}{5}\right) \\ + \sum_{l,m}\left[(3-l){}^{(6)}A_{lm}\xi_1^{l-1} - (l+1)\frac{\kappa_{6:lm}}{\xi_1^{l+2}}\right]Y_l^m. \end{aligned} \quad (\text{H25})$$

Therefore, we obtain

$$\tilde{c}_6 = \frac{45\mu^2}{2(1+p)^2\xi_1^2}, \quad (\text{H26})$$

$$\begin{aligned} \Theta_6 = {}^{(6)}A_2\xi^2P_2(\sin\theta\cos\varphi) + {}^{(6)}A_{22}\xi^2P_2^2(\cos\theta)\cos 2\varphi + {}^{(6)}A_4\xi^4P_4(\sin\theta\cos\varphi) \\ + {}^{(6)}A_5\xi^5P_5(\sin\theta\cos\varphi), \end{aligned} \quad (\text{H27})$$

where

$${}^{(6)}A_2 = \frac{225\mu^2}{28(1+p)^2\xi_1^4}, \quad (\text{H28})$$

$${}^{(6)}A_{22} = \frac{75\mu^2}{8(1+p)\xi_1^4}, \quad (\text{H29})$$

$${}^{(6)}A_4 = \frac{45\mu^2}{14(1+p)^2\xi_1^6}, \quad (\text{H30})$$

$${}^{(6)}A_5 = -\frac{11\mu}{8(1+p)\xi_1^5}. \quad (\text{H31})$$

2. The case of $n = 1$

In this case, since we can express ${}^{(3)}\psi_2 = {}^{(3)}B_2j_2$ and ${}^{(6)}\psi_{lm} = {}^{(6)}B_{lm}$ except for ${}^{(6)}\psi_{22}$ by using the constants ${}^{(3)}B_2$ and ${}^{(6)}B_{lm}$, the boundary conditions can be written as

$$\begin{aligned} \tilde{c}_6 + \frac{\mu}{1+p}P_5(\sin\theta\cos\varphi) - \frac{2\mu}{\xi_1}{}^{(4)}\phi_2(\xi_1)P_2^2(\cos\theta)\cos 2\varphi \\ = \frac{{}^{(3)}B_2j_2(\xi_1)}{\Theta_{0,\xi}(\xi_1)}\left[-\frac{2\mu}{\xi_1(1+p)} + {}^{(3)}B_2\frac{dj_2}{d\xi}(\xi_1) + \frac{3\kappa_{3:2}}{\xi_1^4}\right]\left(\frac{18}{35}P_4(\sin\theta\cos\varphi) + \frac{2}{7}P_2(\sin\theta\cos\varphi) + \frac{1}{5}\right) \\ + \sum_{l,m}\left(-{}^{(6)}B_{lm}j_l(\xi_1) + \frac{\kappa_{6:lm}}{\xi_1^{l+1}}\right)Y_l^m, \end{aligned} \quad (\text{H32})$$

$$\begin{aligned} \frac{5\mu}{\xi_1(1+p)}P_5(\sin\theta\cos\varphi) - \frac{2\mu}{\xi_1}\frac{d^{(4)}\phi_2}{d\xi}(\xi_1)P_2^2(\cos\theta)\cos 2\varphi \\ = \frac{{}^{(3)}B_2j_2(\xi_1)}{\Theta_{0,\xi}(\xi_1)}\left[-\frac{2\mu}{\xi_1^2(1+p)} + \left(\frac{3}{\xi_1^2} + \frac{\xi_1}{2}\frac{d^3\Theta_0}{d\xi^3}(\xi_1)\right){}^{(3)}B_2j_2(\xi_1) + {}^{(3)}B_2\frac{d^2j_2}{d\xi^2}(\xi_1) - \frac{12\kappa_{3:2}}{\xi_1^5}\right] \end{aligned}$$

$$\begin{aligned} & \times \left(\frac{18}{35} P_4(\sin \theta \cos \varphi) + \frac{2}{7} P_2(\sin \theta \cos \varphi) + \frac{1}{5} \right) \\ & + \sum_{l,m} \left[-{}^{(6)}B_{lm} \frac{dj_l}{d\xi}(\xi_1) - (l+1) \frac{\kappa_{6:lm}}{\xi_1^{l+2}} \right] Y_l^m. \end{aligned} \quad (\text{H33})$$

Therefore, we obtain

$$\tilde{c}_6 = \frac{45\mu^2}{2(1+p)^2\xi_1^2}, \quad (\text{H34})$$

$$\begin{aligned} \Theta_6 = & {}^{(6)}B_{2j_2}(\xi) P_2(\sin \theta \cos \varphi) + {}^{(6)}\psi_{22}(\xi) P_2^2(\cos \theta) \cos 2\varphi + {}^{(6)}B_{4j_4}(\xi) P_4(\sin \theta \cos \varphi) \\ & + {}^{(6)}B_{5j_5}(\xi) P_5(\sin \theta \cos \varphi), \end{aligned} \quad (\text{H35})$$

where

$${}^{(6)}B_2 = \frac{225\mu^2}{7(1+p)^2\xi_1^2}, \quad (\text{H36})$$

$${}^{(6)}B_4 = \frac{405\mu^2}{7(1+p)^2(15-\xi_1^2)}, \quad (\text{H37})$$

$${}^{(6)}B_5 = -\frac{11\mu\xi_1^3}{(1+p)(105-10\xi_1^2)}, \quad (\text{H38})$$

$$j_5(\xi) = \frac{1}{\xi^6} \left[(945 - 420\xi^2 + 15\xi^4) \sin \xi - \xi(945 - 105\xi^2 + \xi^4) \cos \xi \right], \quad (\text{H39})$$

and the coefficient of ${}^{(6)}\psi_{22}$ is determined from the equation;

$$3{}^{(6)}\psi_{22}(\xi_1) + \xi_1 \frac{d({}^{(6)}\psi_{22})}{d\xi}(\xi_1) = \frac{2\mu}{\xi_1} \left[3{}^{(4)}\phi_2(\xi_1) + \xi_1 \frac{d({}^{(4)}\phi_2)}{d\xi}(\xi_1) \right]. \quad (\text{H40})$$

3. The case of $n > 0$

In this case, the boundary conditions can be written as

$$\begin{aligned} \tilde{c}_6 + \frac{\mu}{1+p} P_5(\sin \theta \cos \varphi) - \frac{2\mu}{\xi_1} {}^{(4)}\phi_2(\xi_1) P_2^2(\cos \theta) \cos 2\varphi \\ = \frac{{}^{(3)}\psi_2(\xi_1)}{\Theta_{0,\xi}(\xi_1)} \left[-\frac{2\mu}{\xi_1(1+p)} + \frac{d({}^{(3)}\psi_2)}{d\xi}(\xi_1) + \frac{3\kappa_{3:2}}{\xi_1^4} \right] \left(\frac{18}{35} P_4(\sin \theta \cos \varphi) + \frac{2}{7} P_2(\sin \theta \cos \varphi) + \frac{1}{5} \right) \\ + \sum_{l,m} \left[-{}^{(6)}\psi_{lm}(\xi_1) + \frac{\kappa_{6:lm}}{\xi_1^{l+1}} \right] Y_l^m, \end{aligned} \quad (\text{H41})$$

$$\begin{aligned} \frac{5\mu}{\xi_1(1+p)} P_5(\sin \theta \cos \varphi) - \frac{2\mu}{\xi_1} \frac{d({}^{(4)}\phi_2)}{d\xi}(\xi_1) P_2^2(\cos \theta) \cos 2\varphi \\ = \frac{{}^{(3)}\psi_2(\xi_1)}{\Theta_{0,\xi}(\xi_1)} \left[-\frac{2\mu}{\xi_1^2(1+p)} + \left(\frac{3}{\xi_1^2} - \frac{1}{2\Theta_{0,\xi}(\xi_1)} \frac{d^3\Theta_0}{d\xi^3}(\xi_1) \right) {}^{(3)}\psi_2(\xi_1) + \frac{d^2({}^{(3)}\psi_2)}{d\xi^2}(\xi_1) - \frac{12\kappa_{3:2}}{\xi_1^5} \right] \\ \times \left(\frac{18}{35} P_4(\sin \theta \cos \varphi) + \frac{2}{7} P_2(\sin \theta \cos \varphi) + \frac{1}{5} \right) \\ + \sum_{l,m} \left[-\frac{d({}^{(6)}\psi_{lm})}{d\xi}(\xi_1) - (l+1) \frac{\kappa_{6:lm}}{\xi_1^{l+2}} \right] Y_l^m. \end{aligned} \quad (\text{H42})$$

Therefore, we obtain

$$\tilde{c}_6 = -\frac{\xi_1}{10\Theta_{0,\xi}(\xi_1)} n \Theta_0^{n-1}(\xi_1) ({}^{(3)}\psi_2(\xi_1))^2 - {}^{(6)}\psi_0(\xi_1) - \xi_1 \frac{d({}^{(6)}\psi_0)}{d\xi}(\xi_1), \quad (\text{H43})$$

$$\begin{aligned} \Theta_6 = & {}^{(6)}\psi_0(\xi) + {}^{(6)}\psi_2(\xi) P_2(\sin \theta \cos \varphi) + {}^{(6)}\psi_{22}(\xi) P_2^2(\cos \theta) \cos 2\varphi + {}^{(6)}\psi_4(\xi) P_4(\sin \theta \cos \varphi) \\ & + {}^{(6)}\psi_5(\xi) P_5(\sin \theta \cos \varphi), \end{aligned} \quad (\text{H44})$$

where the forms of ${}^{(6)}\psi_i$ are determined by the equations

$$\frac{1}{\xi^2} \frac{d}{d\xi} \left(\xi^2 \frac{d}{d\xi} \right) {}^{(6)}\psi_0 = -n\Theta_0^{n-1} {}^{(6)}\psi_0 - \frac{1}{10} n(n-1) \Theta_0^{n-2} ({}^{(3)}\psi_2)^2, \quad (\text{H45})$$

$$\left[\frac{1}{\xi^2} \frac{d}{d\xi} \left(\xi^2 \frac{d}{d\xi} \right) - \frac{6}{\xi^2} \right] {}^{(6)}\psi_2 = -n\Theta_0^{n-1} {}^{(6)}\psi_2 - \frac{1}{7} n(n-1) \Theta_0^{n-2} ({}^{(3)}\psi_2)^2, \quad (\text{H46})$$

$$\left[\frac{1}{\xi^2} \frac{d}{d\xi} \left(\xi^2 \frac{d}{d\xi} \right) - \frac{6}{\xi^2} \right] {}^{(6)}\psi_{22} = -n\Theta_0^{n-1} {}^{(6)}\psi_{22} + \frac{2\mu}{\xi_1} \left[\frac{1}{\xi^2} \frac{d}{d\xi} \left(\xi^2 \frac{d}{d\xi} \right) - \frac{6}{\xi^2} \right] {}^{(4)}\phi_2, \quad (\text{H47})$$

$$\left[\frac{1}{\xi^2} \frac{d}{d\xi} \left(\xi^2 \frac{d}{d\xi} \right) - \frac{20}{\xi^2} \right] {}^{(6)}\psi_4 = -n\Theta_0^{n-1} {}^{(6)}\psi_4 - \frac{9}{35} n(n-1) \Theta_0^{n-2} ({}^{(3)}\psi_2)^2, \quad (\text{H48})$$

$$\left[\frac{1}{\xi^2} \frac{d}{d\xi} \left(\xi^2 \frac{d}{d\xi} \right) - \frac{30}{\xi^2} \right] {}^{(6)}\psi_5 = -n\Theta_0^{n-1} {}^{(6)}\psi_5, \quad (\text{H49})$$

and the coefficients of ${}^{(6)}\psi_i$ are determined by the boundary conditions;

$$3 {}^{(6)}\psi_2(\xi_1) + \xi_1 \frac{d {}^{(6)}\psi_2}{d\xi}(\xi_1) = -\frac{\xi_1}{7\Theta_{0,\xi}(\xi_1)} n\Theta_0^{n-1}(\xi_1) ({}^{(3)}\psi_2(\xi_1))^2, \quad (\text{H50})$$

$$3 {}^{(6)}\psi_{22}(\xi_1) + \xi_1 \frac{d {}^{(6)}\psi_{22}}{d\xi}(\xi_1) = \frac{2\mu}{\xi_1} \left[3 {}^{(4)}\phi_2(\xi_1) + \xi_1 \frac{d {}^{(4)}\phi_2}{d\xi}(\xi_1) \right], \quad (\text{H51})$$

$$5 {}^{(6)}\psi_4(\xi_1) + \xi_1 \frac{d {}^{(6)}\psi_4}{d\xi}(\xi_1) = -\frac{9\xi_1}{35\Theta_{0,\xi}(\xi_1)} n\Theta_0^{n-1}(\xi_1) ({}^{(3)}\psi_2(\xi_1))^2, \quad (\text{H52})$$

$$6 {}^{(6)}\psi_5(\xi_1) + \xi_1 \frac{d {}^{(6)}\psi_5}{d\xi}(\xi_1) = -\frac{11\mu}{1+p}. \quad (\text{H53})$$

APPENDIX I: THE CHANGE IN THE CENTRAL DENSITY

In this section, we present the expression of the change in the central density at $O(\epsilon^6)$ by using the first tensor virial relation.

First, we divide the first tensor virial relation (2.28),

$$\frac{3}{n} \Pi_1 + \frac{3}{n'} \Pi_2 + (W_{self})_{tot} + (W_{int})_{tot} + 2T_{tot} = 0, \quad (\text{I1})$$

into three parts of $O(\epsilon^0)$, $O(\epsilon)$ and $O(\epsilon^6)$. After that we demand that these three parts should be zero independently.

(a) The 0th order of ϵ :

$$\frac{M_1^2}{2a_0} \left[\frac{5-n}{1+n} \frac{1}{\xi_1^3 (\Theta_{0,\xi}(\xi_1))^2} \int_0^{\xi_1} d\xi \xi^2 \Theta_0^{1+n} - 1 \right] + \frac{M_2^2}{2a'_0} \left[\frac{5-n'}{1+n'} \frac{1}{\xi_1'^3 (\bar{\Theta}_{0,\xi'}(\xi'_1))^2} \int_0^{\xi'_1} d\xi' \xi'^2 \bar{\Theta}_0^{1+n'} - 1 \right] = 0. \quad (\text{I2})$$

If we assume that both of the terms concerned with star 1 and star 2 become zero independently, we have two relations;

$$\frac{5-n}{1+n} \frac{1}{\xi_1^3 (\Theta_{0,\xi}(\xi_1))^2} \int_0^{\xi_1} d\xi \xi^2 \Theta_0^{1+n} = 1, \quad (\text{I3})$$

$$\frac{5-n'}{1+n'} \frac{1}{\xi_1'^3 (\bar{\Theta}_{0,\xi'}(\xi'_1))^2} \int_0^{\xi'_1} d\xi' \xi'^2 \bar{\Theta}_0^{1+n'} = 1. \quad (\text{I4})$$

The above relations are proved using the Lane-Emden equations.

(b) The 1st order of ϵ :

It is clear that this term becomes zero.

(c) The 6th order of ϵ :

By using the relations;

$$\tilde{c}_6 = -\left(\frac{3-n}{2n}\right)\xi_1\Theta_{0,\xi}(\xi_1)\frac{\delta\rho_c}{\rho_c} - {}^{(6)}\psi_0(\xi_1), \quad (I5)$$

$$\tilde{c}'_6 = -\left(\frac{3-n'}{2n'}\right)\xi'_1\bar{\Theta}_{0,\xi'}(\xi'_1)\frac{\delta\rho'_c}{\rho'_c} - {}^{(6)}\bar{\psi}_0(\xi'_1), \quad (I6)$$

$$\mu = -\left(\frac{1+p}{p}\right)\xi_1\Theta_{0,\xi}(\xi_1), \quad (I7)$$

$$\mu' = -(1+p)\xi'_1\bar{\Theta}_{0,\xi'}(\xi'_1), \quad (I8)$$

we can express the equation at order ϵ^6 as

$$\begin{aligned} & \frac{M_1^2}{a_0} \left[-\frac{5-n}{2(1+n)} \frac{1}{\xi_1^3(\Theta_{0,\xi}(\xi_1))^2} \left[\frac{5-n}{2n} \frac{\delta\rho_c}{\rho_c} \int_0^{\xi_1} d\xi \xi^2 \Theta_0^{1+n} - (1+n) \int_0^{\xi_1} d\xi \xi^2 \left\{ \Theta_0^{n(6)}\psi_0 + \frac{1}{10} n \Theta_0^{n-1} ({}^{(3)}\psi_2)^2 \right\} \right] \right. \\ & \quad \left. + \frac{15}{4p} \frac{\bar{F}_{11}}{M_1 a_0^2} - \frac{{}^{(6)}\psi_0(\xi_1)}{2\xi_1\Theta_{0,\xi}(\xi_1)} + \frac{n-1}{4n} \frac{\delta\rho_c}{\rho_c} \right] \\ & + \frac{M_2^2}{a'_0} \left(\frac{a'_0}{a_0} \right)^6 \left[-\frac{5-n'}{2(1+n')} \frac{1}{\xi_1'^3(\bar{\Theta}_{0,\xi'}(\xi'_1))^2} \right. \\ & \quad \times \left[\frac{5-n'}{2n'} \frac{\delta\rho'_c}{\rho'_c} \int_0^{\xi'_1} d\xi' \xi'^2 \bar{\Theta}_0^{1+n'} - (1+n') \int_0^{\xi'_1} d\xi' \xi'^2 \left\{ \bar{\Theta}_0^{n'(6)}\bar{\psi}_0 + \frac{1}{10} n' \bar{\Theta}_0^{n'-1} ({}^{(3)}\bar{\psi}_2)^2 \right\} \right] \\ & \quad \left. + \frac{15p}{4} \frac{\bar{F}'_{11}}{M_2 a_0'^2} - \frac{{}^{(6)}\bar{\psi}_0(\xi'_1)}{2\xi'_1\bar{\Theta}_{0,\xi'}(\xi'_1)} + \frac{n'-1}{4n'} \frac{\delta\rho'_c}{\rho'_c} \right] = 0. \end{aligned} \quad (I9)$$

If we assume that both of the terms concerned with star 1 and star 2 become zero independently, we obtain two relations;

$$\frac{\delta\rho_c}{\rho_c} = \frac{2n}{3-n} \left[\frac{15}{4p} \frac{\bar{F}_{11}}{M_1 a_0^2} - \frac{{}^{(6)}\psi_0(\xi_1)}{2\xi_1\Theta_{0,\xi}(\xi_1)} + \frac{5-n}{2\xi_1^3(\Theta_{0,\xi}(\xi_1))^2} \int_0^{\xi_1} d\xi \xi^2 \left\{ \Theta_0^{n(6)}\psi_0 + \frac{1}{10} n \Theta_0^{n-1} ({}^{(3)}\psi_2)^2 \right\} \right], \quad (I10)$$

$$\frac{\delta\rho'_c}{\rho'_c} = \frac{2n'}{3-n'} \left[\frac{15p}{4} \frac{\bar{F}'_{11}}{M_2 a_0'^2} - \frac{{}^{(6)}\bar{\psi}_0(\xi'_1)}{2\xi'_1\bar{\Theta}_{0,\xi'}(\xi'_1)} + \frac{5-n'}{2\xi_1'^3(\bar{\Theta}_{0,\xi'}(\xi'_1))^2} \int_0^{\xi'_1} d\xi' \xi'^2 \left\{ \bar{\Theta}_0^{n'(6)}\bar{\psi}_0 + \frac{1}{10} n' \bar{\Theta}_0^{n'-1} ({}^{(3)}\bar{\psi}_2)^2 \right\} \right], \quad (I11)$$

where we have used Eqs. (I3) and (I4).

- [1] K. Kuroda *et al.*, in *Proceedings of International Conference on Gravitational Waves: Sources and Detectors*, ed. I. Ciufolini and F. Fidecaro (World Scientific, Singapore, 1997), p.100.
- [2] K. Danzmann, in *Gravitational Wave Experiments*, ed. E. Coccia, G. Pizzella, and F. Ronga (World Scientific, Singapore, 1995), p.100.
- [3] B. Caron *et al.*, in *Proceedings of International Conference on Gravitational Waves: Sources and Detectors*, ed. I. Ciufolini and F. Fidecaro (World Scientific, Singapore, 1997), p.73.
- [4] M. W. Coles, in *Proceedings of International Conference on Gravitational Waves: Sources and Detectors*, ed. I. Ciufolini and F. Fidecaro (World Scientific, Singapore, 1997), p.93.
- [5] K. S. Thorne, in *Relativistic Cosmology, Proc. of the 8th Nishinomiya-Yukawa Memorial Symposium*, ed. M. Sasaki (Tokyo Universal Academy, 1994), p.67.
- [6] For example, K. S. Thorne, in *Proceeding of Snowmass 94 Summer Study on Particle and Nuclear Astrophysics and Cosmology*, ed. E. W. Kolb and R. Peccei (World Scientific, Singapore, 1995) and references cited therein.
- [7] L. Lindblom, *Astrophys. J.* **398**, 569 (1992).
- [8] M. Shibata, *Phys. Rev. D* **60**, 104052 (1999).
- [9] M. Shibata and K. Uryū, *Phys. Rev. D* **61**, 064001 (2000).
- [10] K. Oohara and T. Nakamura, to appear in *Prog. Theor. Phys. Suppl.* **136** (1999), preprint astro-ph/9912085.
- [11] For example, F. A Rasio and S. L. Shapiro, *Class. Quant. Grav.* **16**, R1 (1999) and references cited therein.
- [12] S. Bonazzola, E.ourgoulhon, and J.-A. Marck, *Phys. Rev. D* **56**, 7740 (1997).
- [13] H. Asada, *Phys. Rev. D* **57**, 7292 (1998).

- [14] M. Shibata, Phys. Rev. D **58**, 24012 (1998).
- [15] S. A. Teukolsky, Astrophys. J. **504**, 442 (1998).
- [16] C. S. Kochanek, Astrophys. J. **398**, 234 (1992).
- [17] L. Bildsten and C. Cutler, Astrophys. J. **400**, 175 (1992).
- [18] S. Bonazzola, E. Gourgoulhon, and J.-A. Marck, Phys. Rev. Lett. **82**, 892 (1999).
- [19] S. Bonazzola, E. Gourgoulhon, and J.-A. Marck, to appear in the Proceeding of the 19th Texas Symposium on Relativistic Astrophysics, preprint gr-qc/9903105.
- [20] S. Bonazzola, E. Gourgoulhon, P. Grandclement, and J.-A. Marck, to appear in Prog. Theor. Phys. Suppl. **136** (1999), preprint gr-qc/9909020.
- [21] K. Uryū and Y. Eriguchi, accepted to Phys. Rev. D., preprint gr-qc/9908059.
- [22] J. C. Lombardi, F. A. Rasio, and S. L. Shapiro, Phys. Rev. D **56**, 3416 (1997).
- [23] K. Taniguchi, Prog. Theor. Phys. **101**, 283 (1999).
- [24] K. Uryū and Y. Eriguchi, Mon. Not. R. Astron. Soc., **296**, L1 (1998).
- [25] K. Uryū and Y. Eriguchi, Astrophys. J. Suppl. Ser. **118**, 563 (1998).
- [26] D. Lai, F. A. Rasio, and S. L. Shapiro, Astrophys. J. **420**, 811 (1994).
- [27] P. Marronetti, G. J. Mathews, and J. R. Wilson, Phys. Rev. D **60**, 087301 (1999).
- [28] G. J. Mathews and J. R. Wilson, preprint gr-qc/9911047.
- [29] K. Taniguchi and T. Nakamura, Phys. Rev. Lett. **84**, 581 (2000).
- [30] S. Chandrasekhar, Mon. Not. R. Astron. Soc. **93**, 390 (1933).
- [31] A. Kovetz, Astrphys. J. **154**, 999 (1968).
- [32] D. Lai, F. A. Rasio, and S. L. Shapiro, Astrophys. J. Suppl. Ser. **88**, 205 (1993).
- [33] Private communications with S. Bonazzola, E. Gourgoulhon, and J.-A. Marck on the existence of the slight velocity component along the orbital axis.
- [34] S. Chandrasekhar, *Ellipsoidal Figures of Equilibrium* (Yale University Press, New Heaven, CT, 1969).

Tables

TABLE I. The functions and their derivatives to represent the equilibrium figure for $n = 0.5$. (I)

$n = 0.5$								
ξ	Θ_0	$d\Theta_0/d\xi$	$^{(3)}\psi_2$	$d^{(3)}\psi_2/d\xi$	$^{(4)}\psi_3$	$d^{(4)}\psi_3/d\xi$	$^{(5)}\psi_4$	$d^{(5)}\psi_4/d\xi$
0.00	1.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
0.10	0.9983	-3.332(-2)	5.371(-3)	0.1074	-1.316(-4)	-3.946(-3)	4.003(-6)	1.601(-4)
0.20	0.9933	-6.653(-2)	2.146(-2)	0.2143	-1.052(-3)	-1.576(-2)	6.400(-5)	1.279(-3)
0.30	0.9850	-9.955(-2)	4.820(-2)	0.3203	-3.544(-3)	-3.539(-2)	3.236(-4)	4.311(-3)
0.40	0.9734	-0.1323	8.548(-2)	0.4249	-8.385(-3)	-6.270(-2)	1.021(-3)	1.019(-2)
0.50	0.9586	-0.1646	0.1331	0.5277	-1.634(-2)	-9.755(-2)	2.488(-3)	1.985(-2)
0.60	0.9405	-0.1964	0.1909	0.6280	-2.814(-2)	-0.1397	5.146(-3)	3.416(-2)
0.70	0.9193	-0.2276	0.2586	0.7255	-4.452(-2)	-0.1890	9.505(-3)	5.400(-2)
0.80	0.8950	-0.2581	0.3359	0.8197	-6.617(-2)	-0.2450	1.616(-2)	8.017(-2)
0.90	0.8678	-0.2877	0.4224	0.9099	-9.374(-2)	-0.3075	2.577(-2)	0.1134
1.00	0.8375	-0.3165	0.5178	0.9956	-0.1279	-0.3759	3.910(-2)	0.1545
1.10	0.8045	-0.3441	0.6214	1.076	-0.1691	-0.4498	5.695(-2)	0.2040
1.20	0.7687	-0.3707	0.7328	1.151	-0.2180	-0.5287	8.020(-2)	0.2624
1.30	0.7304	-0.3959	0.8514	1.219	-0.2750	-0.6118	0.1097	0.3302
1.40	0.6896	-0.4197	0.9764	1.280	-0.3405	-0.6985	0.1466	0.4078
1.50	0.6465	-0.4420	1.107	1.333	-0.4148	-0.7879	0.1916	0.4952
1.60	0.6013	-0.4626	1.243	1.377	-0.4981	-0.8789	0.2459	0.5926
1.70	0.5540	-0.4814	1.382	1.411	-0.5906	-0.9705	0.3105	0.6996
1.80	0.5051	-0.4982	1.525	1.433	-0.6922	-1.061	0.3862	0.8158
1.90	0.4545	-0.5128	1.668	1.442	-0.8027	-1.149	0.4739	0.9404
2.00	0.4026	-0.5250	1.813	1.436	-0.9219	-1.233	0.5745	1.072
2.10	0.3496	-0.5346	1.955	1.412	-1.049	-1.310	0.6885	1.209
2.20	0.2958	-0.5413	2.094	1.367	-1.183	-1.376	0.8164	1.349
2.30	0.2414	-0.5448	2.228	1.296	-1.324	-1.427	0.9582	1.487
2.40	0.1869	-0.5447	2.352	1.190	-1.468	-1.456	1.114	1.617
2.50	0.1326	-0.5403	2.464	1.037	-1.614	-1.451	1.281	1.728
2.60	7.904(-2)	-0.5306	2.557	0.8061	-1.756	-1.389	1.458	1.798
2.70	2.674(-2)	-0.5139	2.620	0.3959	-1.887	-1.189	1.637	1.753
2.75270	0.000	-0.5000	2.628	-0.3645	-1.942	-0.6776	1.724	1.368

TABLE II. The functions and their derivatives to represent the equilibrium figure for $n = 1$. (I)

ξ	$n = 1.0$							
	Θ_0	$d\Theta_0/d\xi$	$^{(3)}\psi_2$	$d^{(3)}\psi_2/d\xi$	$^{(4)}\psi_3$	$d^{(4)}\psi_3/d\xi$	$^{(5)}\psi_4$	$d^{(5)}\psi_4/d\xi$
0.00	1.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
0.10	0.9983	-3.330(-2)	3.331(-3)	6.657(-2)	-6.977(-5)	-2.092(-3)	1.831(-6)	7.324(-5)
0.20	0.9933	-6.640(-2)	1.330(-2)	0.1326	-5.573(-4)	-8.347(-3)	2.926(-5)	5.847(-4)
0.30	0.9851	-9.910(-2)	2.981(-2)	0.1974	-1.876(-3)	-1.869(-2)	1.478(-4)	1.967(-3)
0.40	0.9735	-0.1312	5.273(-2)	0.2606	-4.428(-3)	-3.302(-2)	4.656(-4)	4.639(-3)
0.50	0.9589	-0.1625	8.186(-2)	0.3216	-8.606(-3)	-5.116(-2)	1.132(-3)	9.006(-3)
0.60	0.9411	-0.1929	0.1169	0.3797	-1.478(-2)	-7.291(-2)	2.336(-3)	1.544(-2)
0.70	0.9203	-0.2221	0.1577	0.4347	-2.330(-2)	-9.804(-2)	4.302(-3)	2.431(-2)
0.80	0.8967	-0.2500	0.2038	0.4859	-3.449(-2)	-0.1263	7.289(-3)	3.591(-2)
0.90	0.8704	-0.2764	0.2547	0.5329	-4.865(-2)	-0.1572	1.158(-2)	5.054(-2)
1.00	0.8415	-0.3012	0.3102	0.5753	-6.602(-2)	-0.1907	1.750(-2)	6.842(-2)
1.10	0.8102	-0.3242	0.3696	0.6128	-8.684(-2)	-0.2261	2.538(-2)	8.974(-2)
1.20	0.7767	-0.3453	0.4326	0.6450	-0.1113	-0.2632	3.557(-2)	0.1147
1.30	0.7412	-0.3644	0.4984	0.6717	-0.1395	-0.3015	4.844(-2)	0.1432
1.40	0.7039	-0.3814	0.5667	0.6925	-0.1716	-0.3405	6.434(-2)	0.1755
1.50	0.6650	-0.3962	0.6367	0.7074	-0.2076	-0.3798	8.367(-2)	0.2115
1.60	0.6247	-0.4087	0.7080	0.7161	-0.2476	-0.4190	0.1068	0.2511
1.70	0.5833	-0.4189	0.7798	0.7186	-0.2914	-0.4575	0.1340	0.2942
1.80	0.5410	-0.4268	0.8515	0.7148	-0.3390	-0.4949	0.1657	0.3405
1.90	0.4981	-0.4323	0.9225	0.7048	-0.3903	-0.5307	0.2022	0.3898
2.00	0.4546	-0.4354	0.9922	0.6886	-0.4451	-0.5645	0.2438	0.4419
2.10	0.4111	-0.4361	1.060	0.6664	-0.5031	-0.5957	0.2907	0.4963
2.20	0.3675	-0.4345	1.125	0.6382	-0.5642	-0.6240	0.3431	0.5527
2.30	0.3242	-0.4307	1.187	0.6044	-0.6278	-0.6490	0.4013	0.6106
2.40	0.2814	-0.4245	1.246	0.5651	-0.6938	-0.6703	0.4653	0.6695
2.50	0.2394	-0.4162	1.300	0.5207	-0.7618	-0.6875	0.5352	0.7289
2.60	0.1983	-0.4058	1.350	0.4715	-0.8312	-0.7004	0.6110	0.7882
2.70	0.1583	-0.3935	1.394	0.4179	-0.9017	-0.7086	0.6928	0.8468
2.80	0.1196	-0.3792	1.433	0.3604	-0.9728	-0.7119	0.7804	0.9041
2.90	8.250(-2)	-0.3633	1.466	0.2993	-1.044	-0.7100	0.8735	0.9595
3.00	4.704(-2)	-0.3457	1.493	0.2352	-1.115	-0.7030	0.9722	1.012
3.10	1.341(-2)	-0.3266	1.513	0.1686	-1.184	-0.6906	1.076	1.062
3.14159	0.000	-0.3183	1.520	0.1402	-1.213	-0.6838	1.120	1.081

TABLE III. The functions and their derivatives to represent the equilibrium figure for $n = 1.5$. (I)

ξ	$n = 1.5$							
	Θ_0	$d\Theta_0/d\xi$	$^{(3)}\psi_2$	$d^{(3)}\psi_2/d\xi$	$^{(4)}\psi_3$	$d^{(4)}\psi_3/d\xi$	$^{(5)}\psi_4$	$d^{(5)}\psi_4/d\xi$
0.00	1.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
0.10	0.9983	-3.328(-2)	1.994(-3)	3.984(-2)	-3.512(-5)	-1.053(-3)	7.821(-7)	3.127(-5)
0.20	0.9934	-6.627(-2)	7.951(-3)	7.917(-2)	-2.802(-4)	-4.194(-3)	1.249(-5)	2.494(-4)
0.30	0.9851	-9.866(-2)	1.779(-2)	0.1175	-9.419(-4)	-9.372(-3)	6.301(-5)	8.375(-4)
0.40	0.9737	-0.1302	3.140(-2)	0.1543	-2.220(-3)	-1.650(-2)	1.982(-4)	1.971(-3)
0.50	0.9591	-0.1605	4.859(-2)	0.1892	-4.303(-3)	-2.547(-2)	4.810(-4)	3.815(-3)
0.60	0.9416	-0.1895	6.916(-2)	0.2218	-7.370(-3)	-3.613(-2)	9.900(-4)	6.521(-3)
0.70	0.9213	-0.2169	9.286(-2)	0.2517	-1.158(-2)	-4.831(-2)	1.818(-3)	1.022(-2)
0.80	0.8983	-0.2424	0.1194	0.2786	-1.708(-2)	-6.184(-2)	3.071(-3)	1.503(-2)
0.90	0.8728	-0.2659	0.1485	0.3022	-2.398(-2)	-7.650(-2)	4.866(-3)	2.106(-2)
1.00	0.8452	-0.2873	0.1797	0.3225	-3.241(-2)	-9.211(-2)	7.326(-3)	2.836(-2)
1.10	0.8155	-0.3064	0.2129	0.3393	-4.243(-2)	-0.1084	1.058(-2)	3.701(-2)
1.20	0.7840	-0.3231	0.2475	0.3526	-5.411(-2)	-0.1253	1.477(-2)	4.703(-2)
1.30	0.7509	-0.3375	0.2833	0.3623	-6.750(-2)	-0.1424	2.004(-2)	5.845(-2)
1.40	0.7166	-0.3494	0.3198	0.3686	-8.260(-2)	-0.1597	2.651(-2)	7.126(-2)
1.50	0.6811	-0.3590	0.3569	0.3716	-9.943(-2)	-0.1769	3.433(-2)	8.545(-2)
1.60	0.6448	-0.3661	0.3940	0.3716	-0.1180	-0.1939	4.364(-2)	0.1010
1.70	0.6080	-0.3710	0.4311	0.3686	-0.1382	-0.2106	5.458(-2)	0.1179
1.80	0.5707	-0.3737	0.4677	0.3631	-0.1601	-0.2268	6.726(-2)	0.1360
1.90	0.5333	-0.3742	0.5036	0.3553	-0.1835	-0.2424	8.182(-2)	0.1554
2.00	0.4959	-0.3728	0.5387	0.3455	-0.2085	-0.2575	9.838(-2)	0.1760
2.10	0.4588	-0.3696	0.5727	0.3341	-0.2350	-0.2720	0.1171	0.1978
2.20	0.4221	-0.3646	0.6054	0.3214	-0.2629	-0.2859	0.1380	0.2207
2.30	0.3859	-0.3582	0.6369	0.3079	-0.2922	-0.2992	0.1613	0.2448
2.40	0.3505	-0.3504	0.6670	0.2938	-0.3227	-0.3120	0.1870	0.2700
2.50	0.3159	-0.3413	0.6957	0.2795	-0.3546	-0.3244	0.2153	0.2965
2.60	0.2823	-0.3313	0.7229	0.2654	-0.3876	-0.3365	0.2463	0.3241
2.70	0.2497	-0.3204	0.7488	0.2519	-0.4219	-0.3485	0.2802	0.3531
2.80	0.2182	-0.3088	0.7733	0.2393	-0.4573	-0.3606	0.3170	0.3836
2.90	0.1879	-0.2967	0.7967	0.2278	-0.4940	-0.3730	0.3569	0.4157
3.00	0.1589	-0.2843	0.8189	0.2180	-0.5319	-0.3860	0.4002	0.4497
3.10	0.1311	-0.2715	0.8403	0.2102	-0.5712	-0.3999	0.4470	0.4859
3.20	0.1046	-0.2588	0.8611	0.2047	-0.6119	-0.4151	0.4975	0.5246
3.30	7.931(-2)	-0.2460	0.8814	0.2021	-0.6543	-0.4323	0.5520	0.5666
3.40	5.534(-2)	-0.2334	0.9016	0.2031	-0.6985	-0.4519	0.6109	0.6124
3.50	3.262(-2)	-0.2212	0.9221	0.2086	-0.7448	-0.4752	0.6746	0.6633
3.60	1.109(-2)	-0.2094	0.9435	0.2206	-0.7937	-0.5041	0.7438	0.7213
3.65375	0.000	-0.2033	0.9557	0.2318	-0.8213	-0.5240	0.7835	0.7575

TABLE IV. The functions and their derivatives to represent the equilibrium figure for $n = 2$. (I)

ξ	$n = 2.0$							
	Θ_0	$d\Theta_0/d\xi$	$^{(3)}\psi_2$	$d^{(3)}\psi_2/d\xi$	$^{(4)}\psi_3$	$d^{(4)}\psi_3/d\xi$	$^{(5)}\psi_4$	$d^{(5)}\psi_4/d\xi$
0.00	1.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
0.10	0.9983	-3.327(-2)	1.125(-3)	2.247(-2)	-1.631(-5)	-4.890(-4)	3.016(-7)	1.206(-5)
0.20	0.9934	-6.614(-2)	4.481(-3)	4.456(-2)	-1.301(-4)	-1.945(-3)	4.812(-6)	9.607(-5)
0.30	0.9851	-9.822(-2)	1.001(-2)	6.589(-2)	-4.366(-4)	-4.337(-3)	2.425(-5)	3.221(-4)
0.40	0.9738	-0.1292	1.762(-2)	8.613(-2)	-1.027(-3)	-7.612(-3)	7.617(-5)	7.563(-4)
0.50	0.9594	-0.1586	2.719(-2)	0.1050	-1.986(-3)	-1.170(-2)	1.845(-4)	1.460(-3)
0.60	0.9421	-0.1863	3.856(-2)	0.1222	-3.392(-3)	-1.653(-2)	3.789(-4)	2.486(-3)
0.70	0.9222	-0.2119	5.156(-2)	0.1375	-5.313(-3)	-2.199(-2)	6.941(-4)	3.883(-3)
0.80	0.8998	-0.2352	6.599(-2)	0.1508	-7.807(-3)	-2.799(-2)	1.169(-3)	5.689(-3)
0.90	0.8752	-0.2561	8.165(-2)	0.1620	-1.093(-2)	-3.443(-2)	1.847(-3)	7.934(-3)
1.00	0.8487	-0.2745	9.833(-2)	0.1712	-1.470(-2)	-4.120(-2)	2.771(-3)	1.064(-2)
1.10	0.8204	-0.2904	0.1158	0.1782	-1.917(-2)	-4.821(-2)	3.991(-3)	1.383(-2)
1.20	0.7907	-0.3036	0.1339	0.1833	-2.435(-2)	-5.538(-2)	5.553(-3)	1.750(-2)
1.30	0.7598	-0.3142	0.1524	0.1864	-3.025(-2)	-6.262(-2)	7.507(-3)	2.166(-2)
1.40	0.7279	-0.3224	0.1711	0.1879	-3.688(-2)	-6.986(-2)	9.902(-3)	2.632(-2)
1.50	0.6954	-0.3281	0.1899	0.1878	-4.422(-2)	-7.706(-2)	1.279(-2)	3.147(-2)
1.60	0.6624	-0.3316	0.2086	0.1863	-5.228(-2)	-8.417(-2)	1.621(-2)	3.710(-2)
1.70	0.6291	-0.3330	0.2272	0.1838	-6.105(-2)	-9.117(-2)	2.022(-2)	4.323(-2)
1.80	0.5958	-0.3325	0.2454	0.1803	-7.051(-2)	-9.805(-2)	2.487(-2)	4.984(-2)
1.90	0.5627	-0.3302	0.2632	0.1762	-8.066(-2)	-0.1048	3.021(-2)	5.693(-2)
2.00	0.5298	-0.3263	0.2806	0.1715	-9.147(-2)	-0.1115	3.628(-2)	6.453(-2)
2.10	0.4975	-0.3211	0.2975	0.1666	-0.1029	-0.1180	4.313(-2)	7.262(-2)
2.20	0.4656	-0.3147	0.3139	0.1616	-0.1151	-0.1245	5.082(-2)	8.124(-2)
2.30	0.4345	-0.3073	0.3298	0.1567	-0.1279	-0.1310	5.940(-2)	9.039(-2)
2.40	0.4042	-0.2991	0.3452	0.1519	-0.1413	-0.1375	6.892(-2)	0.1001
2.50	0.3747	-0.2902	0.3602	0.1475	-0.1554	-0.1441	7.944(-2)	0.1104
2.60	0.3462	-0.2808	0.3748	0.1435	-0.1701	-0.1509	9.102(-2)	0.1214
2.70	0.3186	-0.2710	0.3889	0.1401	-0.1855	-0.1578	0.1037	0.1330
2.80	0.2920	-0.2610	0.4028	0.1372	-0.2017	-0.1650	0.1177	0.1454
2.90	0.2664	-0.2508	0.4164	0.1351	-0.2185	-0.1724	0.1328	0.1586
3.00	0.2418	-0.2406	0.4298	0.1337	-0.2362	-0.1803	0.1494	0.1726
3.10	0.2183	-0.2305	0.4432	0.1330	-0.2546	-0.1886	0.1674	0.1875
3.20	0.1957	-0.2204	0.4565	0.1332	-0.2739	-0.1973	0.1869	0.2035
3.30	0.1742	-0.2106	0.4698	0.1341	-0.2941	-0.2067	0.2081	0.2205
3.40	0.1536	-0.2010	0.4833	0.1359	-0.3153	-0.2166	0.2311	0.2387
3.50	0.1340	-0.1917	0.4970	0.1386	-0.3374	-0.2273	0.2559	0.2582
3.60	0.1153	-0.1827	0.5111	0.1422	-0.3607	-0.2387	0.2828	0.2791
3.70	9.742(-2)	-0.1740	0.5255	0.1466	-0.3852	-0.2510	0.3118	0.3015
3.80	8.043(-2)	-0.1658	0.5404	0.1519	-0.4110	-0.2641	0.3431	0.3255
3.90	6.425(-2)	-0.1579	0.5559	0.1581	-0.4381	-0.2783	0.3769	0.3513
4.00	4.884(-2)	-0.1504	0.5721	0.1653	-0.4666	-0.2935	0.4134	0.3790
4.10	3.416(-2)	-0.1433	0.5890	0.1733	-0.4968	-0.3098	0.4528	0.4087
4.20	2.016(-2)	-0.1367	0.6068	0.1824	-0.5287	-0.3274	0.4952	0.4407
4.30	6.811(-3)	-0.1304	0.6255	0.1924	-0.5623	-0.3463	0.5410	0.4751
4.35287	0.000	-0.1272	0.6358	0.1980	-0.5809	-0.3569	0.5666	0.4944

TABLE V. The functions and their derivatives to represent the equilibrium figure for $n = 0.5$. (II)

ξ	$n = 0.5$									
	$^{(6)}\psi_0$	$d^{(6)}\psi_0/d\xi$	$^{(6)}\psi_2$	$d^{(6)}\psi_2/d\xi$	$^{(6)}\psi_{22}$	$d^{(6)}\psi_{22}/d\xi$	$^{(6)}\psi_4$	$d^{(6)}\psi_4/d\xi$	$^{(6)}\psi_5$	$d^{(6)}\psi_5/d\xi$
0.00	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
0.10	1.720(-10)	1.032(-8)	5.400(-3)	0.1080	9.847(-3)	0.1969	2.061(-6)	8.245(-5)	-1.312(-7)	-6.558(-6)
0.20	1.104(-8)	3.317(-7)	2.158(-2)	0.2155	3.935(-2)	0.3929	3.299(-5)	6.601(-4)	-4.195(-6)	-1.049(-4)
0.30	1.264(-7)	2.535(-6)	4.846(-2)	0.3220	8.838(-2)	0.5873	1.672(-4)	2.231(-3)	-3.183(-5)	-5.301(-4)
0.40	7.150(-7)	1.078(-5)	8.594(-2)	0.4272	0.1567	0.7792	5.291(-4)	5.299(-3)	-1.339(-4)	-1.672(-3)
0.50	2.752(-6)	3.330(-5)	0.1338	0.5305	0.2441	0.9678	1.294(-3)	1.038(-2)	-4.080(-4)	-4.072(-3)
0.60	8.309(-6)	8.410(-5)	0.1920	0.6315	0.3502	1.152	2.689(-3)	1.799(-2)	-1.013(-3)	-8.419(-3)
0.70	2.123(-5)	1.850(-4)	0.2601	0.7297	0.4744	1.331	4.994(-3)	2.868(-2)	-2.184(-3)	-1.554(-2)
0.80	4.805(-5)	3.684(-4)	0.3378	0.8247	0.6162	1.505	8.547(-3)	4.302(-2)	-4.245(-3)	-2.640(-2)
0.90	9.917(-5)	6.801(-4)	0.4249	0.9159	0.7750	1.671	1.374(-2)	6.160(-2)	-7.624(-3)	-4.207(-2)
1.00	1.904(-4)	1.184(-3)	0.5208	1.003	0.9501	1.829	2.103(-2)	8.508(-2)	-1.286(-2)	-6.377(-2)
1.10	3.453(-4)	1.968(-3)	0.6253	1.085	1.140	1.978	3.094(-2)	0.1141	-2.062(-2)	-9.277(-2)
1.20	5.973(-4)	3.151(-3)	0.7377	1.162	1.345	2.116	4.407(-2)	0.1496	-3.170(-2)	-0.1305
1.30	9.942(-4)	4.895(-3)	0.8575	1.234	1.563	2.243	6.109(-2)	0.1922	-4.705(-2)	-0.1783
1.40	1.602(-3)	7.419(-3)	0.9843	1.299	1.793	2.357	8.279(-2)	0.2432	-6.774(-2)	-0.2376
1.50	2.514(-3)	1.102(-2)	1.117	1.358	2.034	2.456	0.1100	0.3036	-9.501(-2)	-0.3101
1.60	3.856(-3)	1.612(-2)	1.256	1.411	2.284	2.539	0.1439	0.3750	-0.1302	-0.3970
1.70	5.806(-3)	2.329(-2)	1.399	1.456	2.542	2.604	0.1855	0.4594	-0.1749	-0.4997
1.80	8.610(-3)	3.335(-2)	1.547	1.494	2.804	2.648	0.2363	0.5593	-0.2307	-0.6194
1.90	1.261(-2)	4.753(-2)	1.698	1.526	3.070	2.668	0.2980	0.6782	-0.2994	-0.7571
2.00	1.831(-2)	6.766(-2)	1.852	1.551	3.337	2.661	0.3727	0.8214	-0.3828	-0.9134
2.10	2.644(-2)	9.663(-2)	2.008	1.572	3.601	2.621	0.4633	0.9966	-0.4827	-1.088
2.20	3.809(-2)	0.1393	2.166	1.592	3.860	2.544	0.5735	1.217	-0.6010	-1.281
2.30	5.505(-2)	0.2045	2.326	1.620	4.109	2.418	0.7089	1.505	-0.7394	-1.489
2.40	8.032(-2)	0.3100	2.491	1.672	4.342	2.231	0.8782	1.906	-0.8992	-1.708
2.50	0.1197	0.4979	2.663	1.797	4.552	1.956	1.097	2.528	-1.081	-1.929
2.60	0.1867	0.9026	2.857	2.158	4.728	1.538	1.402	3.713	-1.284	-2.130
2.70	0.3305	2.398	3.133	3.903	4.849	0.7890	1.917	7.685	-1.504	-2.243
2.75270	0.6770	∞	3.641	∞	4.868	-0.6130	2.888	∞	-1.620	-1.969

TABLE VI. The functions and their derivatives to represent the equilibrium figure for $n = 1$. (II)

$n = 1.0$										
ξ	${}^{(6)}\psi_0$	$d^{(6)}\psi_0/d\xi$	${}^{(6)}\psi_2$	$d^{(6)}\psi_2/d\xi$	${}^{(6)}\psi_{22}$	$d^{(6)}\psi_{22}/d\xi$	${}^{(6)}\psi_4$	$d^{(6)}\psi_4/d\xi$	${}^{(6)}\psi_5$	$d^{(6)}\psi_5/d\xi$
0.00	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
0.10	0.000	0.000	2.170(-3)	4.336(-2)	4.281(-3)	8.556(-2)	1.193(-6)	4.770(-5)	-5.203(-8)	-2.601(-6)
0.20	0.000	0.000	8.660(-3)	8.635(-2)	1.709(-2)	0.1704	1.906(-5)	3.808(-4)	-1.663(-6)	-4.155(-5)
0.30	0.000	0.000	1.942(-2)	0.1286	3.831(-2)	0.2538	9.627(-5)	1.281(-3)	-1.260(-5)	-2.098(-4)
0.40	0.000	0.000	3.434(-2)	0.1697	6.779(-2)	0.3352	3.033(-4)	3.022(-3)	-5.297(-5)	-6.605(-4)
0.50	0.000	0.000	5.332(-2)	0.2094	0.1053	0.4138	7.374(-4)	5.866(-3)	-1.611(-4)	-1.605(-3)
0.60	0.000	0.000	7.617(-2)	0.2473	0.1504	0.4890	1.521(-3)	1.006(-2)	-3.992(-4)	-3.308(-3)
0.70	0.000	0.000	0.1027	0.2831	0.2029	0.5601	2.802(-3)	1.583(-2)	-8.584(-4)	-6.085(-3)
0.80	0.000	0.000	0.1327	0.3165	0.2623	0.6267	4.748(-3)	2.339(-2)	-1.664(-3)	-1.030(-2)
0.90	0.000	0.000	0.1659	0.3471	0.3281	0.6880	7.546(-3)	3.292(-2)	-2.979(-3)	-1.634(-2)
1.00	0.000	0.000	0.2020	0.3747	0.3997	0.7438	1.140(-2)	4.456(-2)	-5.008(-3)	-2.465(-2)
1.10	0.000	0.000	0.2408	0.3992	0.4766	0.7934	1.653(-2)	5.845(-2)	-8.000(-3)	-3.568(-2)
1.20	0.000	0.000	0.2817	0.4201	0.5582	0.8365	2.317(-2)	7.468(-2)	-1.225(-2)	-4.991(-2)
1.30	0.000	0.000	0.3247	0.4375	0.6437	0.8728	3.155(-2)	9.330(-2)	-1.810(-2)	-6.780(-2)
1.40	0.000	0.000	0.3691	0.4511	0.7325	0.9019	4.191(-2)	0.1143	-2.595(-2)	-8.986(-2)
1.50	0.000	0.000	0.4147	0.4607	0.8238	0.9237	5.450(-2)	0.1378	-3.623(-2)	-0.1165
1.60	0.000	0.000	0.4611	0.4664	0.9170	0.9380	6.954(-2)	0.1636	-4.943(-2)	-0.1483
1.70	0.000	0.000	0.5079	0.4681	1.011	0.9446	8.728(-2)	0.1916	-6.607(-2)	-0.1856
1.80	0.000	0.000	0.5546	0.4656	1.106	0.9436	0.1079	0.2218	-8.673(-2)	-0.2287
1.90	0.000	0.000	0.6009	0.4591	1.200	0.9350	0.1317	0.2539	-0.1120	-0.2781
2.00	0.000	0.000	0.6463	0.4485	1.292	0.9189	0.1588	0.2878	-0.1426	-0.3340
2.10	0.000	0.000	0.6905	0.4340	1.383	0.8955	0.1893	0.3233	-0.1791	-0.3967
2.20	0.000	0.000	0.7330	0.4157	1.471	0.8649	0.2235	0.3600	-0.2221	-0.4663
2.30	0.000	0.000	0.7735	0.3936	1.556	0.8276	0.2614	0.3977	-0.2725	-0.5429
2.40	0.000	0.000	0.8116	0.3681	1.637	0.7838	0.3030	0.4361	-0.3310	-0.6265
2.50	0.000	0.000	0.8470	0.3391	1.713	0.7341	0.3486	0.4748	-0.3981	-0.7170
2.60	0.000	0.000	0.8793	0.3071	1.783	0.6787	0.3980	0.5134	-0.4746	-0.8142
2.70	0.000	0.000	0.9083	0.2722	1.848	0.6184	0.4513	0.5516	-0.5611	-0.9179
2.80	0.000	0.000	0.9337	0.2347	1.907	0.5537	0.5083	0.5889	-0.6584	-1.028
2.90	0.000	0.000	0.9552	0.1949	1.959	0.4851	0.5690	0.6250	-0.7669	-1.143
3.00	0.000	0.000	0.9726	0.1532	2.004	0.4135	0.6332	0.6594	-0.8872	-1.264
3.10	0.000	0.000	0.9858	0.1098	2.041	0.3394	0.7008	0.6917	-1.020	-1.388
3.14159	0.000	0.000	0.9899	9.134(-2)	2.055	0.3081	0.7298	0.7044	-1.079	-1.441

TABLE VII. The functions and their derivatives to represent the equilibrium figure for $n = 1.5$. (II)

ξ	$n = 1.5$									
	${}^{(6)}\psi_0$	$d^{(6)}\psi_0/d\xi$	${}^{(6)}\psi_2$	$d^{(6)}\psi_2/d\xi$	${}^{(6)}\psi_{22}$	$d^{(6)}\psi_{22}/d\xi$	${}^{(6)}\psi_4$	$d^{(6)}\psi_4/d\xi$	${}^{(6)}\psi_5$	$d^{(6)}\psi_5/d\xi$
0.00	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
0.10	-7.107(-11)	-4.264(-9)	9.164(-4)	1.831(-2)	1.737(-3)	3.470(-2)	6.423(-7)	2.568(-5)	-1.894(-8)	-9.468(-7)
0.20	-4.536(-9)	-1.359(-7)	3.654(-3)	3.638(-2)	6.926(-3)	6.898(-2)	1.024(-5)	2.043(-4)	-6.050(-7)	-1.511(-5)
0.30	-5.142(-8)	-1.025(-6)	8.178(-3)	5.399(-2)	1.550(-2)	0.1024	5.152(-5)	6.832(-4)	-4.581(-6)	-7.620(-5)
0.40	-2.870(-7)	-4.283(-6)	1.443(-2)	7.091(-2)	2.737(-2)	0.1346	1.614(-4)	1.599(-3)	-1.923(-5)	-2.395(-4)
0.50	-1.085(-6)	-1.292(-5)	2.233(-2)	8.694(-2)	4.237(-2)	0.1652	3.898(-4)	3.071(-3)	-5.838(-5)	-5.805(-4)
0.60	-3.207(-6)	-3.170(-5)	3.178(-2)	0.1019	6.034(-2)	0.1939	7.976(-4)	5.201(-3)	-1.444(-4)	-1.193(-3)
0.70	-7.988(-6)	-6.741(-5)	4.266(-2)	0.1155	8.107(-2)	0.2203	1.455(-3)	8.065(-3)	-3.098(-4)	-2.188(-3)
0.80	-1.755(-5)	-1.290(-4)	5.484(-2)	0.1278	0.1043	0.2442	2.437(-3)	1.171(-2)	-5.989(-4)	-3.690(-3)
0.90	-3.502(-5)	-2.276(-4)	6.817(-2)	0.1385	0.1298	0.2655	3.824(-3)	1.617(-2)	-1.069(-3)	-5.835(-3)
1.00	-6.476(-5)	-3.767(-4)	8.249(-2)	0.1476	0.1573	0.2840	5.697(-3)	2.142(-2)	-1.792(-3)	-8.767(-3)
1.10	-1.126(-4)	-5.916(-4)	9.763(-2)	0.1550	0.1865	0.2996	8.135(-3)	2.745(-2)	-2.854(-3)	-1.264(-2)
1.20	-1.859(-4)	-8.897(-4)	0.1134	0.1605	0.2172	0.3123	1.121(-2)	3.417(-2)	-4.356(-3)	-1.760(-2)
1.30	-2.939(-4)	-1.290(-3)	0.1297	0.1643	0.2489	0.3222	1.499(-2)	4.152(-2)	-6.416(-3)	-2.382(-2)
1.40	-4.479(-4)	-1.812(-3)	0.1462	0.1664	0.2815	0.3293	1.953(-2)	4.937(-2)	-9.166(-3)	-3.144(-2)
1.50	-6.611(-4)	-2.478(-3)	0.1629	0.1666	0.3147	0.3337	2.488(-2)	5.760(-2)	-1.276(-2)	-4.062(-2)
1.60	-9.490(-4)	-3.310(-3)	0.1795	0.1652	0.3482	0.3357	3.106(-2)	6.607(-2)	-1.735(-2)	-5.151(-2)
1.70	-1.329(-3)	-4.330(-3)	0.1959	0.1621	0.3817	0.3355	3.809(-2)	7.462(-2)	-2.312(-2)	-6.427(-2)
1.80	-1.822(-3)	-5.563(-3)	0.2119	0.1575	0.4152	0.3332	4.598(-2)	8.308(-2)	-3.027(-2)	-7.903(-2)
1.90	-2.450(-3)	-7.032(-3)	0.2273	0.1514	0.4483	0.3293	5.470(-2)	9.127(-2)	-3.900(-2)	-9.594(-2)
2.00	-3.237(-3)	-8.764(-3)	0.2421	0.1440	0.4810	0.3239	6.422(-2)	9.903(-2)	-4.953(-2)	-0.1151
2.10	-4.212(-3)	-1.079(-2)	0.2561	0.1353	0.5131	0.3175	7.448(-2)	0.1062	-6.211(-2)	-0.1368
2.20	-5.405(-3)	-1.313(-2)	0.2691	0.1255	0.5445	0.3103	8.542(-2)	0.1125	-7.698(-2)	-0.1610
2.30	-6.849(-3)	-1.582(-2)	0.2811	0.1147	0.5751	0.3027	9.695(-2)	0.1178	-9.440(-2)	-0.1880
2.40	-8.582(-3)	-1.891(-2)	0.2920	0.1030	0.6050	0.2951	0.1090	0.1220	-0.1147	-0.2179
2.50	-1.065(-2)	-2.243(-2)	0.3017	9.043(-2)	0.6342	0.2877	0.1213	0.1248	-0.1381	-0.2508
2.60	-1.309(-2)	-2.646(-2)	0.3101	7.713(-2)	0.6626	0.2810	0.1338	0.1259	-0.1649	-0.2871
2.70	-1.596(-2)	-3.106(-2)	0.3171	6.312(-2)	0.6904	0.2753	0.1464	0.1252	-0.1956	-0.3269
2.80	-1.932(-2)	-3.634(-2)	0.3227	4.839(-2)	0.7177	0.2710	0.1588	0.1224	-0.2304	-0.3704
2.90	-2.325(-2)	-4.242(-2)	0.3268	3.291(-2)	0.7447	0.2683	0.1708	0.1170	-0.2698	-0.4181
3.00	-2.784(-2)	-4.950(-2)	0.3293	1.656(-2)	0.7714	0.2678	0.1821	0.1086	-0.3142	-0.4703
3.10	-3.319(-2)	-5.784(-2)	0.3300	-9.219(-4)	0.7983	0.2698	0.1924	9.651(-2)	-0.3641	-0.5275
3.20	-3.946(-2)	-6.785(-2)	0.3290	-1.994(-2)	0.8255	0.2747	0.2013	7.969(-2)	-0.4199	-0.5902
3.30	-4.684(-2)	-8.021(-2)	0.3260	-4.126(-2)	0.8533	0.2832	0.2081	5.650(-2)	-0.4823	-0.6593
3.40	-5.562(-2)	-9.609(-2)	0.3206	-6.634(-2)	0.8823	0.2959	0.2123	2.398(-2)	-0.5520	-0.7357
3.50	-6.626(-2)	-0.1181	0.3125	-9.850(-2)	0.9127	0.3140	0.2124	-2.432(-2)	-0.6298	-0.8209
3.60	-7.969(-2)	-0.1548	0.3004	-0.1493	0.9453	0.3395	0.2062	-0.1100	-0.7166	-0.9174
3.65375	-8.910(-2)	-0.2311	0.2908	-0.2555	0.9640	0.3585	0.1977	-0.3006	-0.7674	-0.9761

TABLE VIII. The functions and their derivatives to represent the equilibrium figure for $n = 2$. (II)

ξ	$n = 2.0$									
	${}^{(6)}\psi_0$	$d^{(6)}\psi_0/d\xi$	${}^{(6)}\psi_2$	$d^{(6)}\psi_2/d\xi$	${}^{(6)}\psi_{22}$	$d^{(6)}\psi_{22}/d\xi$	${}^{(6)}\psi_4$	$d^{(6)}\psi_4/d\xi$	${}^{(6)}\psi_5$	$d^{(6)}\psi_5/d\xi$
0.00	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
0.10	-6.032(-11)	-3.618(-9)	3.396(-4)	6.782(-3)	6.308(-4)	1.260(-2)	2.728(-7)	1.090(-5)	-6.086(-9)	-3.042(-7)
0.20	-3.839(-9)	-1.149(-7)	1.353(-3)	1.345(-2)	2.513(-3)	2.499(-2)	4.339(-6)	8.643(-5)	-1.943(-7)	-4.852(-6)
0.30	-4.331(-8)	-8.611(-7)	3.022(-3)	1.989(-2)	5.616(-3)	3.698(-2)	2.175(-5)	2.874(-4)	-1.470(-6)	-2.443(-5)
0.40	-2.401(-7)	-3.564(-6)	5.319(-3)	2.599(-2)	9.890(-3)	4.839(-2)	6.779(-5)	6.672(-4)	-6.161(-6)	-7.665(-5)
0.50	-9.002(-7)	-1.063(-5)	8.206(-3)	3.167(-2)	1.527(-2)	5.905(-2)	1.626(-4)	1.269(-3)	-1.868(-5)	-1.854(-4)
0.60	-2.633(-6)	-2.573(-5)	1.164(-2)	3.683(-2)	2.167(-2)	6.883(-2)	3.300(-4)	2.123(-3)	-4.610(-5)	-3.801(-4)
0.70	-6.478(-6)	-5.384(-5)	1.555(-2)	4.141(-2)	2.900(-2)	7.763(-2)	5.962(-4)	3.246(-3)	-9.870(-5)	-6.950(-4)
0.80	-1.404(-5)	-1.012(-4)	1.990(-2)	4.535(-2)	3.716(-2)	8.536(-2)	9.883(-4)	4.640(-3)	-1.904(-4)	-1.168(-3)
0.90	-2.759(-5)	-1.750(-4)	2.460(-2)	4.862(-2)	4.604(-2)	9.200(-2)	1.533(-3)	6.293(-3)	-3.390(-4)	-1.841(-3)
1.00	-5.018(-5)	-2.832(-4)	2.960(-2)	5.119(-2)	5.553(-2)	9.753(-2)	2.255(-3)	8.181(-3)	-5.668(-4)	-2.757(-3)
1.10	-8.566(-5)	-4.342(-4)	3.482(-2)	5.306(-2)	6.551(-2)	0.1020	3.176(-3)	1.027(-2)	-9.001(-4)	-3.962(-3)
1.20	-1.387(-4)	-6.365(-4)	4.019(-2)	5.424(-2)	7.589(-2)	0.1054	4.313(-3)	1.251(-2)	-1.370(-3)	-5.500(-3)
1.30	-2.149(-4)	-8.980(-4)	4.564(-2)	5.474(-2)	8.656(-2)	0.1079	5.681(-3)	1.485(-2)	-2.013(-3)	-7.420(-3)
1.40	-3.206(-4)	-1.226(-3)	5.111(-2)	5.462(-2)	9.743(-2)	0.1095	7.286(-3)	1.725(-2)	-2.868(-3)	-9.768(-3)
1.50	-4.626(-4)	-1.628(-3)	5.654(-2)	5.389(-2)	0.1084	0.1103	9.130(-3)	1.964(-2)	-3.982(-3)	-1.259(-2)
1.60	-6.487(-4)	-2.108(-3)	6.187(-2)	5.263(-2)	0.1195	0.1105	1.121(-2)	2.198(-2)	-5.404(-3)	-1.594(-2)
1.70	-8.870(-4)	-2.671(-3)	6.705(-2)	5.088(-2)	0.1305	0.1103	1.352(-2)	2.420(-2)	-7.189(-3)	-1.986(-2)
1.80	-1.186(-3)	-3.321(-3)	7.204(-2)	4.870(-2)	0.1415	0.1096	1.605(-2)	2.627(-2)	-9.396(-3)	-2.440(-2)
1.90	-1.554(-3)	-4.059(-3)	7.678(-2)	4.615(-2)	0.1524	0.1087	1.877(-2)	2.815(-2)	-1.209(-2)	-2.961(-2)
2.00	-2.001(-3)	-4.888(-3)	8.125(-2)	4.329(-2)	0.1633	0.1077	2.167(-2)	2.978(-2)	-1.534(-2)	-3.556(-2)
2.10	-2.535(-3)	-5.807(-3)	8.543(-2)	4.017(-2)	0.1740	0.1067	2.472(-2)	3.116(-2)	-1.923(-2)	-4.229(-2)
2.20	-3.165(-3)	-6.817(-3)	8.928(-2)	3.684(-2)	0.1846	0.1059	2.789(-2)	3.224(-2)	-2.383(-2)	-4.986(-2)
2.30	-3.901(-3)	-7.919(-3)	9.279(-2)	3.336(-2)	0.1952	0.1053	3.116(-2)	3.301(-2)	-2.923(-2)	-5.834(-2)
2.40	-4.752(-3)	-9.113(-3)	9.595(-2)	2.976(-2)	0.2057	0.1050	3.448(-2)	3.345(-2)	-3.553(-2)	-6.781(-2)
2.50	-5.727(-3)	-1.040(-2)	9.874(-2)	2.607(-2)	0.2162	0.1051	3.783(-2)	3.354(-2)	-4.283(-2)	-7.833(-2)
2.60	-6.835(-3)	-1.178(-2)	0.1012	2.234(-2)	0.2267	0.1057	4.118(-2)	3.327(-2)	-5.124(-2)	-9.000(-2)
2.70	-8.087(-3)	-1.326(-2)	0.1032	1.857(-2)	0.2373	0.1069	4.448(-2)	3.264(-2)	-6.087(-2)	-0.1029
2.80	-9.491(-3)	-1.484(-2)	0.1049	1.478(-2)	0.2481	0.1087	4.769(-2)	3.161(-2)	-7.186(-2)	-0.1171
2.90	-1.106(-2)	-1.652(-2)	0.1062	1.099(-2)	0.2591	0.1112	5.078(-2)	3.019(-2)	-8.434(-2)	-0.1328
3.00	-1.280(-2)	-1.832(-2)	0.1071	7.195(-3)	0.2704	0.1145	5.372(-2)	2.835(-2)	-9.847(-2)	-0.1500
3.10	-1.473(-2)	-2.024(-2)	0.1076	3.392(-3)	0.2820	0.1185	5.644(-2)	2.608(-2)	-0.1144	-0.1690
3.20	-1.685(-2)	-2.228(-2)	0.1078	-4.267(-4)	0.2941	0.1234	5.892(-2)	2.334(-2)	-0.1323	-0.1897
3.30	-1.919(-2)	-2.447(-2)	0.1075	-4.275(-3)	0.3067	0.1292	6.109(-2)	2.012(-2)	-0.1524	-0.2125
3.40	-2.175(-2)	-2.681(-2)	0.1069	-8.172(-3)	0.3200	0.1360	6.292(-2)	1.637(-2)	-0.1749	-0.2374
3.50	-2.455(-2)	-2.933(-2)	0.1059	-1.214(-2)	0.3339	0.1437	6.435(-2)	1.206(-2)	-0.2000	-0.2647
3.60	-2.762(-2)	-3.203(-2)	0.1045	-1.621(-2)	0.3487	0.1524	6.531(-2)	7.129(-3)	-0.2279	-0.2945
3.70	-3.097(-2)	-3.495(-2)	0.1026	-2.042(-2)	0.3645	0.1621	6.575(-2)	1.528(-3)	-0.2590	-0.3270
3.80	-3.462(-2)	-3.810(-2)	0.1004	-2.480(-2)	0.3812	0.1730	6.559(-2)	-4.813(-3)	-0.2934	-0.3626
3.90	-3.860(-2)	-4.152(-2)	9.766(-2)	-2.940(-2)	0.3991	0.1850	6.476(-2)	-1.197(-2)	-0.3316	-0.4014
4.00	-4.293(-2)	-4.522(-2)	9.448(-2)	-3.427(-2)	0.4182	0.1982	6.317(-2)	-2.004(-2)	-0.3738	-0.4438
4.10	-4.765(-2)	-4.926(-2)	9.080(-2)	-3.945(-2)	0.4388	0.2127	6.072(-2)	-2.911(-2)	-0.4205	-0.4900
4.20	-5.280(-2)	-5.366(-2)	8.658(-2)	-4.502(-2)	0.4608	0.2284	5.731(-2)	-3.930(-2)	-0.4720	-0.5404
4.30	-5.840(-2)	-5.846(-2)	8.178(-2)	-5.103(-2)	0.4845	0.2456	5.282(-2)	-5.074(-2)	-0.5287	-0.5954
4.35287	-6.156(-2)	-6.119(-2)	7.899(-2)	-5.442(-2)	0.4977	0.2552	4.996(-2)	-5.735(-2)	-0.5610	-0.6264

TABLE IX. The velocity potentials and their derivatives for $n = 0.5$.

ξ	$n = 0.5$					
	$^{(4)}\phi_2$	$d^{(4)}\phi_2/d\xi$	$^{(5)}\phi_3$	$d^{(5)}\phi_3/d\xi$	$^{(6)}\phi_4$	$d^{(6)}\phi_4/d\xi$
0.00	0.000	0.000	0.000	0.000	0.000	0.000
0.10	1.646(-3)	3.291(-2)	1.439(-5)	4.324(-4)	-1.148(-7)	-4.623(-6)
0.20	6.583(-3)	6.584(-2)	1.149(-4)	1.725(-3)	-1.828(-6)	-3.662(-5)
0.30	1.481(-2)	9.878(-2)	3.879(-4)	3.881(-3)	-9.248(-6)	-1.234(-4)
0.40	2.634(-2)	0.1318	9.196(-4)	6.901(-3)	-2.923(-5)	-2.926(-4)
0.50	4.117(-2)	0.1648	1.797(-3)	1.079(-2)	-7.139(-5)	-5.717(-4)
0.60	5.930(-2)	0.1978	3.106(-3)	1.555(-2)	-1.481(-4)	-9.888(-4)
0.70	8.074(-2)	0.2310	4.936(-3)	2.119(-2)	-2.746(-4)	-1.572(-3)
0.80	0.1055	0.2642	7.373(-3)	2.771(-2)	-4.689(-4)	-2.350(-3)
0.90	0.1336	0.2975	1.051(-2)	3.511(-2)	-7.519(-4)	-3.351(-3)
1.00	0.1650	0.3309	1.442(-2)	4.341(-2)	-1.147(-3)	-4.605(-3)
1.10	0.1998	0.3644	1.922(-2)	5.262(-2)	-1.682(-3)	-6.141(-3)
1.20	0.2379	0.3981	2.498(-2)	6.275(-2)	-2.386(-3)	-7.991(-3)
1.30	0.2794	0.4318	3.180(-2)	7.379(-2)	-3.292(-3)	-1.019(-2)
1.40	0.3243	0.4658	3.977(-2)	8.578(-2)	-4.435(-3)	-1.276(-2)
1.50	0.3725	0.4999	4.899(-2)	9.872(-2)	-5.856(-3)	-1.574(-2)
1.60	0.4242	0.5343	5.955(-2)	0.1126	-7.597(-3)	-1.916(-2)
1.70	0.4794	0.5688	7.155(-2)	0.1275	-9.705(-3)	-2.307(-2)
1.80	0.5380	0.6036	8.509(-2)	0.1435	-1.223(-2)	-2.749(-2)
1.90	0.6001	0.6386	0.1003	0.1604	-1.522(-2)	-3.246(-2)
2.00	0.6658	0.6740	0.1172	0.1784	-1.874(-2)	-3.804(-2)
2.10	0.7349	0.7097	0.1360	0.1975	-2.285(-2)	-4.425(-2)
2.20	0.8077	0.7458	0.1567	0.2178	-2.761(-2)	-5.115(-2)
2.30	0.8841	0.7823	0.1796	0.2392	-3.310(-2)	-5.880(-2)
2.40	0.9642	0.8192	0.2046	0.2618	-3.940(-2)	-6.723(-2)
2.50	1.048	0.8568	0.2320	0.2858	-4.658(-2)	-7.653(-2)
2.60	1.136	0.8949	0.2618	0.3111	-5.473(-2)	-8.676(-2)
2.70	1.227	0.9339	0.2943	0.3380	-6.396(-2)	-9.802(-2)
2.75270	1.277	0.9548	0.3125	0.3528	-6.930(-2)	-0.1044

TABLE X. The velocity potentials and their derivatives for $n = 1$.

ξ	$n = 1.0$					
	$^{(4)}\phi_2$	$d^{(4)}\phi_2/d\xi$	$^{(5)}\phi_3$	$d^{(5)}\phi_3/d\xi$	$^{(6)}\phi_4$	$d^{(6)}\phi_4/d\xi$
0.00	0.000	0.000	0.000	0.000	0.000	0.000
0.10	9.819(-4)	1.964(-2)	7.695(-6)	2.313(-4)	-5.453(-8)	-2.195(-6)
0.20	3.929(-3)	3.930(-2)	6.150(-5)	9.233(-4)	-8.684(-7)	-1.740(-5)
0.30	8.844(-3)	5.901(-2)	2.077(-4)	2.079(-3)	-4.396(-6)	-5.871(-5)
0.40	1.573(-2)	7.878(-2)	4.927(-4)	3.702(-3)	-1.391(-5)	-1.394(-4)
0.50	2.460(-2)	9.863(-2)	9.636(-4)	5.797(-3)	-3.401(-5)	-2.728(-4)
0.60	3.546(-2)	0.1186	1.668(-3)	8.371(-3)	-7.068(-5)	-4.729(-4)
0.70	4.832(-2)	0.1387	2.654(-3)	1.143(-2)	-1.313(-4)	-7.539(-4)
0.80	6.320(-2)	0.1589	3.971(-3)	1.499(-2)	-2.246(-4)	-1.130(-3)
0.90	8.012(-2)	0.1794	5.668(-3)	1.905(-2)	-3.610(-4)	-1.618(-3)
1.00	9.908(-2)	0.2000	7.799(-3)	2.364(-2)	-5.523(-4)	-2.232(-3)
1.10	0.1201	0.2209	1.041(-2)	2.876(-2)	-8.121(-4)	-2.989(-3)
1.20	0.1433	0.2420	1.357(-2)	3.444(-2)	-1.155(-3)	-3.908(-3)
1.30	0.1685	0.2634	1.732(-2)	4.069(-2)	-1.600(-3)	-5.007(-3)
1.40	0.1960	0.2851	2.173(-2)	4.753(-2)	-2.164(-3)	-6.306(-3)
1.50	0.2256	0.3072	2.685(-2)	5.499(-2)	-2.868(-3)	-7.827(-3)
1.60	0.2574	0.3296	3.275(-2)	6.310(-2)	-3.737(-3)	-9.591(-3)
1.70	0.2915	0.3525	3.949(-2)	7.188(-2)	-4.795(-3)	-1.162(-2)
1.80	0.3279	0.3758	4.714(-2)	8.136(-2)	-6.072(-3)	-1.395(-2)
1.90	0.3667	0.3996	5.579(-2)	9.159(-2)	-7.597(-3)	-1.661(-2)
2.00	0.4079	0.4240	6.549(-2)	0.1026	-9.405(-3)	-1.961(-2)
2.10	0.4515	0.4490	7.633(-2)	0.1145	-1.153(-2)	-2.301(-2)
2.20	0.4977	0.4745	8.841(-2)	0.1272	-1.402(-2)	-2.684(-2)
2.30	0.5464	0.5008	0.1018	0.1408	-1.692(-2)	-3.113(-2)
2.40	0.5979	0.5279	0.1166	0.1555	-2.026(-2)	-3.593(-2)
2.50	0.6520	0.5558	0.1329	0.1712	-2.412(-2)	-4.130(-2)
2.60	0.7090	0.5845	0.1509	0.1880	-2.854(-2)	-4.728(-2)
2.70	0.7690	0.6143	0.1706	0.2061	-3.360(-2)	-5.394(-2)
2.80	0.8319	0.6451	0.1922	0.2255	-3.936(-2)	-6.135(-2)
2.90	0.8980	0.6771	0.2157	0.2463	-4.590(-2)	-6.957(-2)
3.00	0.9674	0.7104	0.2415	0.2687	-5.330(-2)	-7.871(-2)
3.10	1.040	0.7450	0.2695	0.2927	-6.167(-2)	-8.884(-2)
3.14159	1.071	0.7599	0.2819	0.3032	-6.546(-2)	-9.337(-2)

TABLE XI. The velocity potentials and their derivatives for $n = 1.5$.

ξ	$n = 1.5$					
	$^{(4)}\phi_2$	$d^{(4)}\phi_2/d\xi$	$^{(5)}\phi_3$	$d^{(5)}\phi_3/d\xi$	$^{(6)}\phi_4$	$d^{(6)}\phi_4/d\xi$
0.00	0.000	0.000	0.000	0.000	0.000	0.000
0.10	5.484(-4)	1.097(-2)	3.754(-6)	1.129(-4)	-2.308(-8)	-9.293(-7)
0.20	2.195(-3)	2.196(-2)	3.001(-5)	4.507(-4)	-3.677(-7)	-7.372(-6)
0.30	4.943(-3)	3.301(-2)	1.014(-4)	1.016(-3)	-1.863(-6)	-2.490(-5)
0.40	8.798(-3)	4.412(-2)	2.409(-4)	1.812(-3)	-5.902(-6)	-5.923(-5)
0.50	1.377(-2)	5.533(-2)	4.717(-4)	2.844(-3)	-1.446(-5)	-1.162(-4)
0.60	1.987(-2)	6.666(-2)	8.177(-4)	4.118(-3)	-3.010(-5)	-2.021(-4)
0.70	2.711(-2)	7.813(-2)	1.303(-3)	5.641(-3)	-5.603(-5)	-3.232(-4)
0.80	3.550(-2)	8.978(-2)	1.954(-3)	7.422(-3)	-9.613(-5)	-4.865(-4)
0.90	4.507(-2)	0.1016	2.797(-3)	9.473(-3)	-1.550(-4)	-6.993(-4)
1.00	5.583(-2)	0.1137	3.858(-3)	1.181(-2)	-2.379(-4)	-9.695(-4)
1.10	6.782(-2)	0.1260	5.168(-3)	1.443(-2)	-3.510(-4)	-1.306(-3)
1.20	8.105(-2)	0.1386	6.755(-3)	1.737(-2)	-5.015(-4)	-1.717(-3)
1.30	9.555(-2)	0.1515	8.653(-3)	2.064(-2)	-6.972(-4)	-2.213(-3)
1.40	0.1114	0.1648	1.089(-2)	2.425(-2)	-9.473(-4)	-2.806(-3)
1.50	0.1285	0.1784	1.352(-2)	2.824(-2)	-1.262(-3)	-3.506(-3)
1.60	0.1471	0.1925	1.655(-2)	3.262(-2)	-1.653(-3)	-4.328(-3)
1.70	0.1670	0.2070	2.005(-2)	3.741(-2)	-2.132(-3)	-5.285(-3)
1.80	0.1885	0.2219	2.405(-2)	4.266(-2)	-2.715(-3)	-6.394(-3)
1.90	0.2114	0.2374	2.860(-2)	4.838(-2)	-3.416(-3)	-7.671(-3)
2.00	0.2360	0.2535	3.375(-2)	5.462(-2)	-4.255(-3)	-9.136(-3)
2.10	0.2621	0.2701	3.954(-2)	6.140(-2)	-5.250(-3)	-1.081(-2)
2.20	0.2900	0.2874	4.605(-2)	6.878(-2)	-6.425(-3)	-1.271(-2)
2.30	0.3196	0.3054	5.332(-2)	7.679(-2)	-7.802(-3)	-1.487(-2)
2.40	0.3511	0.3241	6.143(-2)	8.547(-2)	-9.409(-3)	-1.732(-2)
2.50	0.3845	0.3436	7.044(-2)	9.488(-2)	-1.128(-2)	-2.008(-2)
2.60	0.4199	0.3639	8.043(-2)	0.1051	-1.344(-2)	-2.318(-2)
2.70	0.4573	0.3851	9.148(-2)	0.1161	-1.592(-2)	-2.666(-2)
2.80	0.4969	0.4073	0.1037	0.1280	-1.878(-2)	-3.057(-2)
2.90	0.5388	0.4305	0.1171	0.1409	-2.205(-2)	-3.493(-2)
3.00	0.5830	0.4547	0.1319	0.1548	-2.579(-2)	-3.981(-2)
3.10	0.6298	0.4801	0.1481	0.1698	-3.003(-2)	-4.523(-2)
3.20	0.6791	0.5066	0.1659	0.1860	-3.485(-2)	-5.127(-2)
3.30	0.7311	0.5343	0.1854	0.2035	-4.031(-2)	-5.797(-2)
3.40	0.7860	0.5634	0.2066	0.2223	-4.647(-2)	-6.539(-2)
3.50	0.8438	0.5938	0.2299	0.2424	-5.341(-2)	-7.360(-2)
3.60	0.9048	0.6256	0.2552	0.2641	-6.122(-2)	-8.266(-2)
3.65375	0.9389	0.6433	0.2697	0.2764	-6.580(-2)	-8.790(-2)

TABLE XII. The velocity potentials and their derivatives for $n = 2$.

ξ	$n = 2.0$					
	$^{(4)}\phi_2$	$d^{(4)}\phi_2/d\xi$	$^{(5)}\phi_3$	$d^{(5)}\phi_3/d\xi$	$^{(6)}\phi_4$	$d^{(6)}\phi_4/d\xi$
0.00	0.000	0.000	0.000	0.000	0.000	0.000
0.10	2.801(-4)	5.604(-3)	1.632(-6)	4.909(-5)	-8.494(-9)	-3.420(-7)
0.20	1.121(-3)	1.123(-2)	1.306(-5)	1.962(-4)	-1.354(-7)	-2.716(-6)
0.30	2.527(-3)	1.689(-2)	4.416(-5)	4.430(-4)	-6.869(-7)	-9.190(-6)
0.40	4.502(-3)	2.261(-2)	1.050(-4)	7.917(-4)	-2.179(-6)	-2.190(-5)
0.50	7.052(-3)	2.842(-2)	2.060(-4)	1.246(-3)	-5.347(-6)	-4.310(-5)
0.60	1.019(-2)	3.432(-2)	3.577(-4)	1.809(-3)	-1.116(-5)	-7.519(-5)
0.70	1.392(-2)	4.034(-2)	5.715(-4)	2.487(-3)	-2.083(-5)	-1.207(-4)
0.80	1.826(-2)	4.650(-2)	8.591(-4)	3.286(-3)	-3.584(-5)	-1.825(-4)
0.90	2.323(-2)	5.283(-2)	1.233(-3)	4.212(-3)	-5.797(-5)	-2.637(-4)
1.00	2.883(-2)	5.933(-2)	1.706(-3)	5.276(-3)	-8.933(-5)	-3.675(-4)
1.10	3.510(-2)	6.605(-2)	2.293(-3)	6.485(-3)	-1.324(-4)	-4.978(-4)
1.20	4.205(-2)	7.300(-2)	3.008(-3)	7.850(-3)	-1.899(-4)	-6.587(-4)
1.30	4.971(-2)	8.020(-2)	3.868(-3)	9.384(-3)	-2.653(-4)	-8.548(-4)
1.40	5.810(-2)	8.767(-2)	4.891(-3)	1.110(-2)	-3.622(-4)	-1.091(-3)
1.50	6.725(-2)	9.545(-2)	6.095(-3)	1.301(-2)	-4.850(-4)	-1.373(-3)
1.60	7.720(-2)	0.1036	7.500(-3)	1.513(-2)	-6.386(-4)	-1.708(-3)
1.70	8.798(-2)	0.1120	9.128(-3)	1.748(-2)	-8.285(-4)	-2.101(-3)
1.80	9.962(-2)	0.1209	1.100(-2)	2.008(-2)	-1.061(-3)	-2.561(-3)
1.90	0.1122	0.1301	1.315(-2)	2.294(-2)	-1.343(-3)	-3.097(-3)
2.00	0.1257	0.1398	1.560(-2)	2.609(-2)	-1.683(-3)	-3.717(-3)
2.10	0.1401	0.1500	1.838(-2)	2.955(-2)	-2.090(-3)	-4.432(-3)
2.20	0.1557	0.1607	2.152(-2)	3.335(-2)	-2.573(-3)	-5.252(-3)
2.30	0.1723	0.1718	2.506(-2)	3.751(-2)	-3.144(-3)	-6.192(-3)
2.40	0.1901	0.1836	2.904(-2)	4.206(-2)	-3.816(-3)	-7.263(-3)
2.50	0.2090	0.1960	3.349(-2)	4.703(-2)	-4.602(-3)	-8.480(-3)
2.60	0.2293	0.2089	3.846(-2)	5.245(-2)	-5.517(-3)	-9.860(-3)
2.70	0.2508	0.2226	4.399(-2)	5.835(-2)	-6.579(-3)	-1.142(-2)
2.80	0.2738	0.2369	5.014(-2)	6.477(-2)	-7.808(-3)	-1.318(-2)
2.90	0.2983	0.2520	5.697(-2)	7.175(-2)	-9.222(-3)	-1.515(-2)
3.00	0.3242	0.2678	6.451(-2)	7.932(-2)	-1.085(-2)	-1.737(-2)
3.10	0.3518	0.2844	7.285(-2)	8.753(-2)	-1.270(-2)	-1.985(-2)
3.20	0.3811	0.3019	8.204(-2)	9.641(-2)	-1.483(-2)	-2.261(-2)
3.30	0.4122	0.3202	9.216(-2)	0.1060	-1.724(-2)	-2.570(-2)
3.40	0.4452	0.3395	0.1033	0.1164	-1.998(-2)	-2.912(-2)
3.50	0.4802	0.3596	0.1155	0.1275	-2.307(-2)	-3.291(-2)
3.60	0.5172	0.3808	0.1288	0.1395	-2.657(-2)	-3.711(-2)
3.70	0.5564	0.4029	0.1434	0.1524	-3.051(-2)	-4.174(-2)
3.80	0.5978	0.4261	0.1593	0.1663	-3.494(-2)	-4.685(-2)
3.90	0.6416	0.4503	0.1767	0.1812	-3.990(-2)	-5.246(-2)
4.00	0.6879	0.4756	0.1956	0.1971	-4.544(-2)	-5.861(-2)
4.10	0.7368	0.5020	0.2161	0.2140	-5.164(-2)	-6.535(-2)
4.20	0.7883	0.5296	0.2384	0.2322	-5.853(-2)	-7.270(-2)
4.30	0.8427	0.5583	0.2626	0.2515	-6.620(-2)	-8.073(-2)
4.35287	0.8726	0.5739	0.2762	0.2622	-7.059(-2)	-8.525(-2)

TABLE XIII. The normalized radii for spherical stars, the orbital separations at the contact point, the energies, the reduced quadrupole moments, and the changes in the central density for $n = 0.5, 1, 1.5$ and 2 .

n	ξ_1	R_c/a_0	$\tilde{E}_{self}$	$\tilde{E}_{point}$	$\tilde{E}_{quad}$	$\tilde{I}_{11}/(Ma_0^2)$	$\delta\rho_c/\rho_c$
0.5	2.75270	2.496	-1.111	-0.5000	1.962	0.6691	2.069
1.0	3.14159	2.426	-1.000	-0.5000	1.040	0.3465	2.280
1.5	3.65375	2.384	-0.8571	-0.5000	0.5749	0.1915	2.158
2.0	4.35287	2.358	-0.6667	-0.5000	0.2958	9.861(-2)	1.924

TABLE XIV. The individual energies for $n = 0.5, 1, 1.5$ and 2 .

n	$\tilde{\Pi}_{self}$	$\tilde{\Pi}_{quad}$	$(\tilde{W}_{self})_{self}$	$(\tilde{W}_{self})_{quad}$	$(\tilde{W}_{int})_{point}$	$(\tilde{W}_{int})_{quad}$	$\tilde{T}_{point}$	$\tilde{T}_{quad}$
0.5	0.2222	-0.9946	-1.333	1.953	-1.000	-2.007	0.5000	3.011
1.0	0.5000	-1.300	-1.500	1.819	-1.000	-1.040	0.5000	1.559
1.5	0.8571	-1.437	-1.714	1.724	-1.000	-0.5746	0.5000	0.8620
2.0	1.333	-1.479	-2.000	1.627	-1.000	-0.2958	0.5000	0.4437

TABLE XV. The total energy, the total angular momentum and the orbital angular velocity along the equilibrium sequences of the binary system.

R/a_0	$E/(M^2/a_0)$	$J/(M^3 a_0)^{1/2}$	$\Omega/(\pi \rho_0)^{1/2}$
$n = 0.5$			
7.0	-1.183	1.871	8.819(-2)
6.5	-1.188	1.803	9.857(-2)
6.0	-1.194	1.733	0.1112
5.5	-1.202	1.659	0.1267
5.0	-1.211	1.583	0.1462
4.5	-1.222	1.502	0.1713
4.0	-1.236	1.418	0.2047
3.5	-1.253	1.330	0.2508
3.0	-1.275	1.240	0.3181
2.5	-1.303	1.152	0.4257
2.496	-1.303	1.151	0.4269
$n = 1$			
7.0	-1.071	1.871	8.818(-2)
6.5	-1.077	1.803	9.855(-2)
6.0	-1.083	1.732	0.1111
5.5	-1.091	1.659	0.1266
5.0	-1.100	1.582	0.1461
4.5	-1.111	1.501	0.1712
4.0	-1.125	1.416	0.2044
3.5	-1.142	1.327	0.2501
3.0	-1.165	1.233	0.3163
2.5	-1.196	1.136	0.4197
2.426	-1.201	1.122	0.4400
$n = 1.5$			
7.0	-0.9286	1.871	8.818(-2)
6.5	-0.9341	1.803	9.855(-2)
6.0	-0.9405	1.732	0.1111
5.5	-0.9480	1.659	0.1266
5.0	-0.9571	1.582	0.1461
4.5	-0.9682	1.501	0.1711
4.0	-0.9820	1.415	0.2043
3.5	-0.9997	1.325	0.2498
3.0	-1.023	1.229	0.3154
2.5	-1.055	1.128	0.4167
2.384	-1.064	1.104	0.4487
$n = 2$			
7.0	-0.7381	1.871	8.818(-2)
6.5	-0.7436	1.803	9.854(-2)
6.0	-0.7500	1.732	0.1111
5.5	-0.7576	1.658	0.1266
5.0	-0.7666	1.581	0.1461
4.5	-0.7777	1.500	0.1711
4.0	-0.7916	1.415	0.2042
3.5	-0.8094	1.324	0.2496
3.0	-0.8329	1.227	0.3148
2.5	-0.8654	1.123	0.4150
2.358	-0.8770	1.092	0.4539

Figures




FIG. 1. (a) The total energy and (b) the total angular momentum as functions of the orbital separation, and (c) the orbital angular velocity as functions of the total angular momentum for the polytropic index $n = 0.5$. Solid line denotes our results. Filled circles and open triangles are the results of Lai, Rasio and Shapiro (1994) and Uryū and Eriguchi (1998).




FIG. 2. (a) The total energy and (b) the total angular momentum as functions of the orbital separation, and (c) the orbital angular velocity as functions of the total angular momentum for the polytropic index $n = 1$. Solid line denotes our results. Filled circles and open triangles are the results of Lai, Rasio and Shapiro (1994) and Uryū and Eriguchi (1998).




FIG. 3. (a) The total energy and (b) the total angular momentum as functions of the orbital separation, and (c) the orbital angular velocity as functions of the total angular momentum for the polytropic index $n = 1.5$. Solid line denotes our results. Filled circles and open triangles are the results of Lai, Rasio and Shapiro (1994) and Uryū and Eriguchi (1998).




FIG. 4. (a) The total energy and (b) the total angular momentum as functions of the orbital separation, and (c) the orbital angular velocity as functions of the total angular momentum for the polytropic index $n = 2$. Solid line denotes our results.




FIG. 5. The relative errors for the functions presented the deformations of the figure for the polytropic index $n = 1$. Thick solid, dotted, dashed, long-dashed, dot-dashed and thin solid lines denote the relative errors for $^{(3)}\psi_2$, $^{(4)}\psi_3$, $^{(5)}\psi_4$, $^{(6)}\psi_2$, $^{(6)}\psi_4$ and $^{(6)}\psi_5$, respectively.